

1-1-2011

# Contemporary and historical separation of transequatorial migration between genetically distinct seabird populations

M.J. Rayner  
*University of Auckland*

M.E. Hauber  
*Hunter University*


T.E. Steeves  
*University of Canterbury*

H.A. Lawrence  
*Manaaki Whenua Landcare Research*

D.R. Thompson  
*National Institute of Water & Atmospheric Research Ltd*

*See next page for additional authors*

Follow this and additional works at: [http://scholarworks.sjsu.edu/biol\\_pub](http://scholarworks.sjsu.edu/biol_pub)

 Part of the [Biology Commons](#), and the [Marine Biology Commons](#)

## Recommended Citation

M.J. Rayner, M.E. Hauber, T.E. Steeves, H.A. Lawrence, D.R. Thompson, P.M. Sagar, S.J. Bury, R.A. Phillips, T.J. Landers, L. Ranjard, and Scott A. Shaffer. "Contemporary and historical separation of transequatorial migration between genetically distinct seabird populations" *Nature Communications* (2011): 332. DOI: 10.1038/ncomms1330

---

**Authors**

M.J. Rayner, M.E. Hauber, T.E. Steeves, H.A. Lawrence, D.R. Thompson, P.M. Sagar, S.J. Bury, R.A. Phillips, T.J. Landers, L.Ranjard, and Scott A. Shaffer

1 Contemporary and historical separation of transequatorial migration between  
2 two genetically-distinct seabird populations

3

4 Matt J. Rayner<sup>1, 2 \*</sup>, Mark E. Hauber<sup>3</sup>, Tammy E. Steeves<sup>4</sup>, Hayley A. Lawrence<sup>5</sup>, David R.  
5 Thompson<sup>6</sup>, Paul M. Sagar<sup>7</sup>, Sarah J. Bury<sup>6</sup>, Todd J. Landers<sup>8, 2</sup>, Richard A. Phillips<sup>9</sup>, Louis  
6 Ranjard<sup>2</sup>, & Scott A. Shaffer<sup>10</sup>

7

8 <sup>1</sup> National Institute of Water & Atmospheric Research Ltd, Private Bag 99940,  
9 Auckland, New Zealand.

10 <sup>2</sup> School of Biological Sciences, University of Auckland, Private Bag 92019, Auckland, New Zealand.

11 <sup>3</sup> Department of Psychology, Hunter College of the City University of New York, 695 Park Avenue, New  
12 York NY 10065, USA.

13 <sup>4</sup> School of Biological Sciences, University of Canterbury, Canterbury, Private Bag 4800, Christchurch,  
14 New Zealand.

15 <sup>5</sup> Manaaki Whenua Landcare Research, Private Bag 92170, Auckland, New Zealand.

16 <sup>6</sup> National Institute of Water & Atmospheric Research Ltd, Private Bag 14901, Wellington, New Zealand.

17 <sup>7</sup> National Institute of Water & Atmospheric Research Ltd, P O Box 8602, Christchurch, New Zealand.

18 <sup>8</sup> Auckland Museum, Private Bag 92018, Auckland, New Zealand.

19 <sup>9</sup> British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road,  
20 Cambridge CB3 0ET, UK.

21 <sup>10</sup> Department of Biological Sciences, San Jose State University, San Jose, 95192-0100, USA.

22

23 \* [m.rayner@niwa.co.nz](mailto:m.rayner@niwa.co.nz)

24

25

26

27

28

29

30

31 **Abstract**

32 Pelagic seabirds are highly mobile, reducing the likelihood of allopatric speciation  
33 where disruption of gene flow between populations is caused by physically  
34 insurmountable, extrinsic barriers. Segregation during the non-breeding season appears  
35 to provide an intrinsic barrier to gene flow among seabird populations that otherwise  
36 occupy nearby or overlapping regions during breeding, but how this is achieved remains  
37 unclear. Here we show that the two genetically distinct populations of Cook's petrel  
38 (*Pterodroma cookie*) exhibit transequatorial separation of non-breeding ranges at  
39 contemporary (ca. 2-3 yrs) and historical (ca. 100 yrs) time scales. Segregation during  
40 the non-breeding season *per se* appears an unlikely barrier gene flow. Instead we  
41 provide evidence that habitat specialisation during the non-breeding season is associated  
42 with breeding asynchrony which, in conjunction with philopatry, restricts gene flow.  
43 Habitat specialisation during breeding and non-breeding likely promotes evolutionary  
44 divergence between these two populations via local adaptation.

45

46

47

48

49

50

51

52

53

54

55

56 **Introduction**

57 Divergent migratory behaviours to and from breeding sites have led to the disruption of  
58 gene flow among populations in many species <sup>1,2</sup>, contributing to reproductive isolation  
59 under the classic model of allopatric speciation <sup>3</sup>. In terrestrial environments, genetic  
60 differentiation among migratory populations is frequently paralleled by extrinsic (e.g.  
61 mountain ranges) and intrinsic (e.g. timing of dispersal, inherited migratory direction,  
62 host use by parasites) barriers that restrict gene flow and facilitate divergence via  
63 genetic drift and/or selection <sup>4-6</sup>. However, the nature of the extrinsic and intrinsic  
64 barriers that disrupt gene flow among populations of marine animals have only recently  
65 begun to be investigated <sup>7,8</sup> and remain poorly known for many highly mobile taxa <sup>8,9</sup>.

66

67 Seabirds undertake the longest known migrations on Earth, routinely crossing  
68 hemispheres between breeding and non-breeding habitat within ocean basins <sup>10,11</sup>. As a  
69 result of this extreme mobility, seabirds experience few apparent physical barriers to  
70 dispersal. Thus, seabirds can potentially visit and ultimately breed on islands thousands  
71 of kilometres from their natal colony, which may contribute to a lack of genetic  
72 structure observed in several species <sup>12-15</sup>. Yet other seabird species show surprisingly  
73 high levels of population genetic structure <sup>8,16-18</sup> and intrinsic barriers to gene flow  
74 appear to play an important role in the evolution of seabird diversity <sup>4,18-22</sup>. For example  
75 given the strong tendency for seabirds to return to their natal site to breed, natal  
76 philopatry has been proposed as such a barrier to gene flow <sup>19</sup>. However given that not  
77 all seabirds exhibit strong genetic structure it is unlikely to be the sole driver.

78

79

80 A recent meta-analysis suggests that segregation of non-breeding distributions may be a  
81 key determinant of genetic structure among seabird populations <sup>8</sup>, but how occupying  
82 disjunct non-breeding distributions can restrict gene flow remains unclear. Although  
83 advances in tracking technology have improved our ability to record the long-distance  
84 movements of seabirds at sea <sup>20</sup>, few studies have examined the non-breeding  
85 segregation amongst seabird populations that occupy nearby or overlapping regions  
86 during the breeding season <sup>11,21</sup>. Accordingly, the role of seasonal movements in  
87 shaping the genetic structure of seabird populations, and the persistence of these  
88 movement patterns at the timescale of generations, remain unknown. Such information  
89 is vital for providing insights into behavioural and ecological mechanisms underlying  
90 the diversification of highly mobile taxa, including those within oceanic environments.

91  
92 *Pterodroma cookii* (Cook's petrel; Procellariiformes: Procellariidae Gray 1843), are  
93 pelagic seabirds that breed exclusively on two islands at the northern and southern  
94 extremes of their prehistoric range within the New Zealand archipelago <sup>22</sup>: Little Barrier  
95 Island (LBI) and Codfish Island (CDF), respectively <sup>23,24</sup>. When breeding, *P. cookii*  
96 range within the south-west Pacific and Tasman Sea, with overlapping foraging ranges  
97 centred around their respective breeding colonies <sup>25</sup>. Shipboard observations indicate  
98 that after breeding, *P. cookii* are trans-equatorial and trans-Pacific migrants, with  
99 sightings concentrated in the North Pacific (North-Pacific Convergence and centrally  
100 along Baja California) and in the South Pacific (Humboldt Current) Oceans <sup>26</sup>. On the  
101 basis of body size and plumage characteristics, in 1929 Murphy <sup>27</sup> distinguished  
102 between smaller *P. c. cookii* that were collected in the North Pacific and assigned to the  
103 breeding population at LBI, and the larger *P. c. orientalis*, of unknown breeding origin,  
104 collected south of the Equator, off the coast of Peru. Subsequently, Falla <sup>28</sup> observed

105 that the plumage and morphological characteristics of *P. cookii orientalis* were  
106 consistent with juveniles of *P. cookii cookii* from the breeding population at CDF that  
107 was discovered in 1934<sup>24</sup>. In addition to differences in body size, body mass, and the  
108 potential for non-breeding birds to occupy disjunct habitats<sup>5,26,29</sup>, birds on LBI and  
109 CDF exhibit a one-month asynchrony in breeding phenology<sup>5</sup> and are genetically  
110 distinct<sup>6</sup>. These combined observations strongly suggest that gene flow between the  
111 two extant *P. cookii* populations is highly restricted. Thus, *P. cookii* is an ideal study  
112 species to explicitly examine whether and how occupying disjunct non-breeding  
113 distributions can restrict gene flow among seabird populations.

114

115 Here we combined geolocator-based tracking with an isotopic ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) and  
116 genetic (mitochondrial cytochrome Oxidase subunit 1) comparison of modern breeding  
117 birds, and historical specimens (i.e. museum skins), to test the hypothesis of  
118 transequatorial separation of the two *P. cookii* breeding populations at varying temporal  
119 scales. Specifically, we tested the predictions that contemporary *P. cookii* populations,  
120 tracked over a complete annual cycle, would exhibit divergent non-breeding  
121 distributions, habitat use, and foraging patterns; this divergence would be consistent  
122 between contemporary cycles (ca. 2-3 years) as revealed by stable isotope analyses; the  
123 stable isotope signatures of historical specimens of unknown breeding provenance  
124 collected ca. 100 years ago from the two non-breeding oceanic regions used by  
125 contemporary populations, would match those of modern samples, and that genetic  
126 analysis would clearly assign origin of breeding population for these museum  
127 specimens and confirm divergence in transequatorial migration of the *P. cookii*  
128 populations over historical timescales. This combination of tracking, museum-based  
129 investigation, and elemental and molecular analyses presents a novel opportunity to

130 evaluate how differences in contemporary and historical migratory behaviour can  
131 contribute to the diversification of a highly mobile marine species.

132

## 133 **Results**

### 134 **Population movements**

135 Consistent with the prediction that contemporary tracked *P. cookii* would exhibit  
136 divergent non-breeding distributions, our analysis reveals that birds tracked between  
137 consecutive breeding seasons using light-based geolocation loggers (hereafter called  
138 loggers) between 2007 and 2009 from LBI (n = 11: female (♀) = 5, male (♂) = 6) and  
139 CDF (n = 11: ♀ = 6, ♂ = 5) exhibited transequatorial separation of their non-breeding  
140 habitats within the Pacific Ocean (Fig. 1, Supplementary Fig. S1). This contrasts with  
141 previous studies of other transequatorial migrants, including related Procellariiform  
142 seabirds, which showed substantial overlaps in space use and mixing during the  
143 nonbreeding period of individuals that originated from different breeding populations  
144 <sup>10,11,29</sup>. Post-breeding LBI *P. cookii* (tracked for  $389 \pm 49$  SD days) completed an anti-  
145 clockwise migration of  $48,037 \pm 7953$  SD km within the North and South Pacific  
146 Ocean. Birds tracked from LBI moved east, then north across the equator to reach core  
147 non-breeding distributions within the California and North-Pacific currents  
148 (approximately 35°N) in  $34 \pm 8$  SD days. Pre-breeding migration returning to New  
149 Zealand waters was completed in  $20 \pm 5$  SD days (Fig. 1) on a direct southwest route.  
150 In contrast, post-breeding CDF *P. cookii* (tracked for  $450 \pm 3$  SD days) migrated within  
151 the South Pacific Ocean ( $37,813 \pm 6920$  SD km) moving east then north within the  
152 Humboldt Current to reach core non-breeding distributions off the Peruvian Coast  
153 (approximately 15° S) in  $19 \pm 5$  SD days. Pre-breeding migration returning to New  
154 Zealand was again completed in  $20 \pm 5$  SD days through a south-western corridor, north



155 of the region traversed during post-breeding movements. Consistent with their breeding  
156 timetables, the migration schedule of both *P. cookii* populations was asynchronous; LBI  
157 birds commenced post- and pre-breeding migrations approximately 1 month before  
158 CDF birds (post-breeding 11 March  $\pm$  12 SD days vs. 8 April  $\pm$  15 SD days ( $t_{(18)} = -$   
159 4.67,  $p < 0.0001$ ) ; pre-breeding 5 Sept  $\pm$  10 SD days vs. 20 Oct  $\pm$  6 SD days ( $t_{(15)} = -$   
160 11.85,  $p < 0.0001$ )).

161

162 During initial migration east towards South America, the routes taken by *P. cookii* from  
163 LBI and CDF overlapped, followed a similar direction to those of post-breeding sooty  
164 shearwaters (*Puffinus griseus*)<sup>11</sup> and Westland petrels (*Procellaria westlandica*)<sup>30</sup>, but  
165 not Australasian gannets (*Morus serrator*), departing from New Zealand nesting  
166 colonies<sup>31</sup>. Subsequently, LBI birds moved north-west across the equator and  
167 eventually returned on south-westerly trajectories to New Zealand along routes that  
168 were directionally similar to sooty shearwaters<sup>11</sup>, flesh-footed shearwaters (*Puffinus*  
169 *carneipes*)<sup>32</sup> and bar-tailed godwits (*Limosa lapponica baueri*) departing from on or off  
170 North America<sup>33</sup>. These observations highlight the existence of an important cross-taxa  
171 avian migration corridor within the Pacific Ocean between approximately 170° E and  
172 160° W.

173

#### 174 **Isotopic and habitat segregation**

175 *P. cookii* moult during the non-breeding period<sup>26,34</sup>, and hence incorporate local  
176 isotopic dietary signals in their new plumage<sup>35</sup>. Stable isotope ratios of C ( $\delta^{13}\text{C}$ ) and N  
177 ( $\delta^{15}\text{N}$ ) in particular provide an indication of carbon source (benthic vs. pelagic, inshore  
178 vs. offshore, and information on water mass) and trophic level of prey, respectively<sup>35</sup>,  
179 and comparisons thus can indicate geographic and/or dietary segregation<sup>36,37</sup>.

180 Consistent with the prediction that geographic divergence would be consistent between  
181 the *P. cookii* populations at a contemporary timescale (ca. 2-3 years), there was no  
182 significant difference in the isotope signatures of *P. cookii* body feathers collected in  
183 2006 and 2008 from LBI (2006 untracked n = 20, 2008 tracked n = 11)( $\delta^{13}\text{C}$ : Z = 0.83,  
184 p = 0.41;  $\delta^{15}\text{N}$ : Z = 1.70, p = 0.10) or CDF (2006 untracked n = 20, 2008 tracked n = 9)(  
185  $\delta^{13}\text{C}$ : Z = 1.34, p = 0.19;  $\delta^{15}\text{N}$ : Z = 1.79, p = 0.07)(Fig. 2) suggesting that, currently,  
186 within each breeding population, birds forage within similar oceanic regions and on  
187 prey of a similar trophic level in successive years.

188

189 In support of the prediction of a match between the stable isotope signatures of  
190 historical and contemporary samples sharing the same non-breeding provenance,  
191 variation in the  $\delta^{13}\text{C}$  signatures of feathers from modern (LBI: 2006 and 2008  
192 combined; CDF: 2006 and 2008 combined) and historical specimens collected within  
193 the core non-breeding habitats of birds tracked from LBI and CDF, respectively  
194 (samples collected in 1905 at 22.42°N, 112.67°W, North Pacific Ocean, in the  
195 California Current region off Baja California, hereafter called BC, and in 1913 at ca.  
196 11°S, 79°W, South Pacific Ocean, in the Humboldt Current, hereafter called HC;  
197 Supplementary Tables S1 and S2) suggest that birds have used population-specific  
198 oceanic regions, with similar carbon signatures, at historical time scales (Fig. 1) (F =  
199 81.32, df = 2, p < 0.001). The mean  $\delta^{13}\text{C}$  value of LBI *P. cookii* ( $-17.18 \pm 0.70$  SD ‰)  
200 was not significantly different from that of historical samples from BC ( $-16.72 \pm 0.51$   
201 SD ‰) (t = 2.085, p = 0.250), but was significantly lower than that of modern samples  
202 from CDF ( $-14.69 \pm 0.58$  ‰) (t = 14.92, p < 0.01), and historical samples from HC ( $-$   
203  $14.79 \pm 0.73$  ‰) (t = 9.292, p < 0.001) (Fig. 2). The mean  $\delta^{13}\text{C}$  value of modern CDF  
204 samples was not significantly different from that of historical samples collected within

205 the HC ( $t = 0.42$   $p = 1.00$ ) but was significantly higher than historical samples collected  
206 from BC ( $t = 8.94$   $p < 0.001$ ) (Fig. 2). These results suggest consistent differences in  
207 carbon isotopic signals between the North Pacific Convergence and South Pacific  
208 Humboldt Current systems, and support our initial prediction.

209

210 The mean  $\delta^{15}\text{N}$  value of modern LBI *P. cookii* feathers ( $15.37 \pm 0.95\text{‰}$ ) was not  
211 significantly different from modern CDF feathers ( $16.05 \pm 1.29\text{‰}$ ) ( $t = 3.69$ ,  $p = 0.16$ )  
212 and the only difference in mean  $\delta^{15}\text{N}$  value between modern and historical feathers was  
213 observed in modern samples from LBI ( $15.37 \pm 0.95\text{‰}$ ) and historical samples from HC  
214 ( $16.76 \pm 1.42\text{‰}$ ) ( $t = 3.53$ ,  $p < 0.01$ ) (Fig. 2). The most likely explanation for these  
215 results is that modern *P. cookii* feed on prey at similar trophic levels, despite occupying  
216 different oceanic regions (as suggested by tracking and carbon isotope data).  
217 Accordingly, the environmental characteristics of core non-breeding habitats exploited  
218 by modern LBI and CDF *P. cookii* were significantly different, with LBI *P. cookii*  
219 foraging over deeper ( $4930 \pm 581$  SD m vs.  $3330 \pm 314$  SD m,  $Z = -3.97$ ,  $p < 0.0001$ ),  
220 warmer ( $20.11 \pm 1.97$  SD °C vs.  $17.29 \pm 1.59$  SD °C,  $Z = 3.12$ ,  $p < 0.01$ ) and less  
221 productive waters ( $0.16 \pm 0.10$  SD mg Chl *a*  $\text{m}^{-3}$  vs.  $0.73 \pm 0.18$  SD mg Chl *a*  $\text{m}^{-3}$ ,  $Z = -$   
222  $3.91$ ,  $p < 0.0001$ ) than their CDF conspecifics (Fig. 3). At the intrapopulation level,  
223 comparisons of environmental characteristics for presence and absence data indicated  
224 that LBI *P. cookii* occupied less productive waters ( $0.16 \pm 0.10$  mg Chl *a*  $\text{m}^{-3}$  vs.  $0.28 \pm$   
225  $0.17$  mg Chl *a*  $\text{m}^{-3}$ ,  $Z = -2.20$ ,  $p = 0.03$ ) and CDF *P. cookii* more productive waters ( $0.73$   
226  $\pm 0.18$  mg Chl *a*  $\text{m}^{-3}$  vs.  $0.47 \pm 0.12$  mg Chl *a*  $\text{m}^{-3}$ ,  $Z = 2.79$ ,  $p < 0.01$ ) than those  
227 available within their respective non-breeding core habitats (Fig. 3).

228

229

## 230 **Genetic divergence**

231 Consistent with prediction that genetic analysis would confirm divergence in  
232 transequatorial migration of the *P. cookii* populations over historical timescales, all nine  
233 historical BC *P. cookii* skins had mitochondrial *cytochrome c oxidase subunit 1* (COI)  
234 haplotypes that were identical to those of the modern LBI population<sup>38</sup> (Fig. 1). Seven  
235 of ten historical HC *P. cookii* skins shared the same haplotype as that of modern CDF *P.*  
236 *cookii*<sup>38</sup> and each of the three remaining skins had a novel haplotype (Fig. 1). We  
237 previously identified a single nucleotide polymorphism (SNP; at site 156) that  
238 differentiates modern LBI and CDF *P. cookii*<sup>38</sup>. All haplotypes sequenced from  
239 historical HC *P. cookii* skins in this study share the same diagnostic SNP as modern  
240 CDF birds, including the three novel haplotypes (Fig. 1, Supplementary Figure S2,  
241 Supplementary Tables S1, S2 and S3). These results support the previous predictions of  
242 contemporary and historical transequatorial separation of the extant *P. cookii*  
243 populations, revealed by our tracking and isotopic analyses, and suggests that these two  
244 populations have been genetically structured for a minimum of 100 years.

245

## 246 **Discussion**

247 Intrinsic factors argued to restrict gene flow between seabird populations include natal  
248 philopatry<sup>19</sup>, divergent breeding timetables<sup>39,40</sup> and differences in breeding and non-  
249 breeding distributions<sup>8,12,17</sup>, though how these interact is poorly understood. Our  
250 combined results suggest that the use of population specific non-breeding habitats plays  
251 an interactive role in the restriction of gene flow, as both a pre-mating and post-mating  
252 barrier. In regards to the former, migration routes of differing lengths or directions, are  
253 widely recognised pre-mating barriers in migrant terrestrial bird species<sup>5,41,42</sup>. In *P.*  
254 *cookii* differences in the duration of migratory movement or non-breeding distribution

255 seem to underlie the pronounced breeding asynchrony between the populations.  
256 Moreover, regional differences in the timing of peak primary productivity, or the onset  
257 of gonad development based upon differences in day length and temperature shifts <sup>43</sup>,  
258 could amplify asynchronies in breeding timetables beyond those mediated by  
259 differences in route or non-breeding residency times alone. Thus, it is not segregation  
260 during the non-breeding season *per se*. that represents a pre-mating barrier to gene flow.  
261 Rather, it is the interaction between differing non-breeding distributions and divergent  
262 breeding timetables, combined with high natal philopatry, which restricts gene flow.

263

264 In regards to the latter, adult Cook's petrel desert their chicks up to two weeks prior to  
265 fledging and, as seen in many migratory species <sup>5,41</sup>, chicks presumably follow an  
266 inherited migratory direction and timing <sup>5,42</sup>. Studies of migrant terrestrial bird species  
267 indicate that the inheritance of suboptimal migration routes during secondary contact  
268 represents a post-mating barrier to gene flow <sup>10, 40</sup>. Although secondary contact is  
269 presumably rare in *P. cookii*, the inheritance of an alternative or intermediate migratory  
270 strategy would place individuals at a selective disadvantage if it led to the occupancy of  
271 suboptimal non-breeding grounds or breeding timetables that were out of synchrony  
272 with the rest of the population <sup>10, 40</sup>.

273

274 Habitat specialisation during both the breeding and non-breeding season is also likely to  
275 promote divergence among the two extant *P. cookii* populations via local adaptation.  
276 For example, when breeding, *P. cookii* exhibit divergent, although overlapping  
277 distributions and diets in response to regional differences in oceanography <sup>25 (current study)</sup>.  
278 Moreover, the populations differ in body mass <sup>22</sup> with smaller LBI birds foraging in  
279 warmer low-latitude waters compared to larger CDF conspecifics <sup>25</sup>. That individuals

280 from these two populations also occupy oceanic waters with similarly divergent  
281 characteristics during the non-breeding period (i.e., LBI birds occupy warmer less  
282 productive waters than CDF birds; Fig. 1) suggests population-specific adaptation to  
283 differing marine habitats, possibly linked to body size. Indeed, differences in seabird  
284 habitat use related to body size have been observed in other studies, particularly in  
285 species with pronounced sexual size dimorphism<sup>44-46</sup>. It remains uncertain whether  
286 body size differences between populations of *P. cookii* are a cause or consequence of  
287 divergent habitat use. Regardless, our data support and extend local adaptation  
288 hypotheses by suggesting that conspecific seabird populations can become tied to  
289 particular marine habitats during the breeding<sup>18,39,47</sup> and non-breeding periods (current  
290 study), presenting opportunities for population divergence.

291

292 Overall, this study demonstrates the use of integrative techniques to track space use,  
293 behaviour, foraging, and breeding to demonstrate clear spatial segregation of non-  
294 breeding distribution between two populations which, mediated by its influence on  
295 breeding phenology, appears to represent an intrinsic barrier to gene flow, and has lead  
296 to local adaptation in a highly mobile seabird.

297

## 298 **Methods**

### 299 **Tracking and environmental data**

300 *P. cookii* breeding on LBI and CDF were equipped with light-based geolocation loggers  
301 (British Antarctic Survey, Mk14) in November-December 2007 (LBI, 36° 11'S, 175°  
302 04'E, n = 13; CDF, 46° 11'S, 167° 38'E, n = 14) during early incubation. Loggers were  
303 deployed on birds of known breeding history using published<sup>25</sup> with the total package  
304 weighing < 1% of body mass. The sex of all birds were previously determined using

305 molecular methods <sup>25,48</sup>. The following breeding season, 24 (89%) birds (LBI n = 12  
306 and CDF n = 12) returned to breed and were recaptured in January - March 2009 at their  
307 same breeding burrows, and loggers were removed. Two loggers (LBI n = 1 and CDF n  
308 = 1) failed to download resulting in 11 datasets from each population for processing and  
309 analyses.

310

311 Light data from the loggers were processed using Multitrace software (Jensen Software  
312 Systems) and locations ( $2 \text{ d}^{-1}$ ) were estimated with an expected mean accuracy  $\pm$  SD of  
313  $186 \pm 114 \text{ km}$  <sup>49</sup>. Sunrise and sunset times were identified based on light curve  
314 thresholds, with latitude calculated from day/night length and longitude calculated from  
315 the time of local midday/midnight relative to Greenwich Mean Time. As a result of day  
316 length uniformity around the equinoxes, clearly erroneous locations occurring 3-4  
317 weeks either side of the equinox were excluded. Furthermore, points that involved  
318 movements of  $> 1600 \text{ km}$  in one day <sup>50</sup>, those with interruptions to light curves around  
319 sunset and sunrise, or that were clearly outside of the known or possible range for *P.*  
320 *cookii*, were removed from the data set.

321

322 Totals of 7,886 and 9,082 locations were obtained for LBI and CDF *P. cookii*  
323 respectively, of which 9.7% and 10.6% were excluded after filtering. Filtered locations  
324 were then used to estimate year-round utilisation distribution (UD) kernels for each  
325 population following methods detailed in ref <sup>51</sup>. In brief, 2-D Gaussian kernel densities  
326 were estimated using custom routines created in MatLab <sup>51</sup> (The Mathworks, Natick,  
327 MA, USA). Kernels were calculated using a Lambert Cylindrical Equal Area projection  
328 on an 80 km grid with grid cells normalized for bird effort by dividing each cell by the  
329 number of birds contributing the locations within a cell. The kernel smoothing

330 parameter (h) was estimated using an adaptive method <sup>52</sup> to estimate an optimal local  
331 value, following ref. <sup>53</sup>. Based published methods <sup>25</sup>, a 1,000 km buffer was used  
332 around each colony to define breeding habitat, and the 80% contour of UD kernels,  
333 calculated individually, were used to define the core non-breeding distributions for each  
334 bird <sup>50</sup>. Dates of the first and last locations to enter and exit the core non-breeding  
335 habitats were used to define migration timing for each individual. Individuals' migration  
336 distances were calculated by summing the point to point distances travelled during post-  
337 breeding and pre-breeding movements between these core areas.

338

339 Satellite-derived remotely-sensed environmental data were used to contrast the core  
340 habitats of *P. cookii* from LBI and CDF during the non-breeding period (presence  
341 dataset) as well as those available, but not used, within each population-specific region  
342 (absence dataset). For the presence dataset, environmental data were extracted for  
343 logger locations falling within the 20% UD for each tracked bird using a 1° longitude by  
344 2° latitude grid centred (the approximate error of geolocation estimates) on the date and  
345 coordinates of the location <sup>51</sup>. For the absence dataset, environmental data were  
346 extracted from ten randomly selected locations, derived within a 1,000 km radius buffer  
347 of each logger location and centred on the corresponding logger location date.  
348 Environmental data extracted for the random locations were averaged to give a mean  
349 absence estimate to contrast presence values. Remotely-sensed environmental data  
350 were obtained from NOAA's Environmental Research Division  
351 (<http://coastwatch.pfel.noaa.gov/thredds/catalog.html>) including 5-day composites of  
352 chlorophyll *a* (Chl *a*) concentration (mg Chl m<sup>-3</sup>) at a spatial resolution of 0.1°  
353 resolution <sup>54</sup> and 8-day composites sea surface temperature (SST) in °C with a spatial  
354 resolution of 0.1° <sup>55</sup>. Bathymetric data were obtained from the ETOPO2 dataset <sup>56</sup>.



355

### 356 **Stable isotopes**

357 To contrast population-specific dietary signals during the non-breeding period, we  
358 conducted stable isotope analyses of C ( $\delta^{13}\text{C}$ ) and N ( $\delta^{15}\text{N}$ ) ratios in body feathers  
359 collected from modern LBI and CDF *P. cookii* and museum specimens held at the  
360 Californian Academy of Sciences (n = 9) and American Museum of Natural History (n  
361 = 10). Single body feathers were collected from breeding *P. cookii* in 2006 and from  
362 tracked birds upon their recapture in 2009, with no resampling of birds between years.  
363 Modern and historical feathers were cleaned with 70% ethanol, then washed in distilled  
364 water to remove contaminants, dried in at 50°C and cut into fine fragments using  
365 scissors. Stable isotope analyses of  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ , using a subsample (approximately  
366 0.7 mg) of each homogenised feather, were carried out on a DeltaPlus (Thermo-  
367 Finnigan) continuous flow isotope ratio mass spectrometer using protocols outlined by  
368 ref<sup>25</sup>. Measurement precision of the isotope analysis was 0.29‰ for  $\delta^{15}\text{N}$  and 0.24‰  
369 for  $\delta^{13}\text{C}$ . To account for the Seuss effect, the temporal biasing of the atmospheric CO<sub>2</sub>  
370 pool to more negative  $\delta^{13}\text{C}$  values as a result of the burning of fossil CO<sub>2</sub>,  $\delta^{13}\text{C}$  data  
371 from historical feathers were normalised by subtracting a year-specific factor  $\delta^{13}\text{C} = -1$   
372  $+ 1.1^{(2009-\text{year} (1905 \text{ or } 1913))*0.027}$  following ref<sup>57</sup>.

373

### 374 **Genetic analyses**

375 Toe pad skin samples were collected from the same historical Cook's petrel specimens  
376 from which feathers were sampled for stable isotope analyses (Supplementary Table S1  
377 and S2). DNA extraction and polymerase chain reaction (PCR) set up was performed in  
378 a physically isolated dedicated ancient DNA laboratory. Contamination was monitored  
379 using extraction and PCR negative controls and genomic DNA was isolated as per ref

380 <sup>58</sup>. A 375 base pair (bp) region of the mitochondrial *cytochrome c oxidase subunit 1*  
381 gene (COI) was amplified using two sets of primers that amplified two overlapping  
382 fragments: AWCF1 and AWCintR2 <sup>58</sup>; LCRintF2 5'-TCATAATTGGGGGATTTGGA-  
383 3' (designed using Primer3 software <sup>59</sup>) and AWCintR3 <sup>58</sup>. This region corresponds to  
384 the first 375 bp of the 677 bp fragment sequenced by ref <sup>38</sup>. PCR products were purified  
385 using a QIAquick PCR Purification kit (Qiagen) then sequenced bidirectionally from  
386 independent PCRs <sup>58</sup>. Sequences were concatenated and aligned using Sequencher  
387 version 4.8 (Gene Codes), deposited in NCBI GenBank (accession numbers HQ263645  
388 to HQ263663) and compared with those in ref <sup>38</sup>.

389

#### 390 **Statistical analyses**

391 Habitat data extractions, processing and analyses, were conducted using Matlab. Non-  
392 parametric Mann-Whitney U tests were used to test for differences between presence  
393 and absence habitat data for the extracted environmental parameters at an intra-  
394 population level, and for the presence data at an inter-population level. Mean  
395 environmental parameter values for individual birds were used as the sampling unit.  
396 Mann-Whitney U tests were initially used to compare stable isotope values between  
397 years (2006 and 2009) for LBI and CDF. Isotope data ( $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ ) were  
398 subsequently pooled for each population and permutation-based non-parametric  
399 ANOVA, with pair-wise comparisons using the Bonferroni correction method,  
400 conducted to compare isotope values between individuals from the breeding  
401 populations (LBI and CDF) and museum skin specimens (CAS 1905 and AMNH 1913)  
402 using the programme PERMANOVA. Sex was initially included as a factor in all  
403 analyses, but excluded as a result of non-significant differences.

404

405 **Acknowledgements**

406 We thank B. Dunphy and New Zealand Department of Conservation (S. McInnes, L.  
407 Witwell, R. Renwick, P. McClelland) for field assistance, M. Flannery (California  
408 Academy of Sciences) and P. Sweet and T. Trombone (The American Museum of  
409 Natural History) for access to museum specimens, C. Millar and S. Patel of assistance  
410 with genetic analyses, J. Brown for assistance with isotopic laboratory support, and W.  
411 Rayner for support during this research. Artwork provided by Chris Gaskin. This work  
412 was supported by the Hauturu Little Barrier Island Supporters Trust (ASB Bank  
413 Community Trust grant), the New Zealand Tertiary Education Commission and  
414 Foundation for Research Science and Technology. This research represents a  
415 contribution to the BAS Ecosystems Programme.

416

417 **Author Contributions**

418 M.J.R, D.R.T, P.M.S, M.E.H, S.A.S, and T.E.S designed the research. M.J.R, T.J.L,  
419 R.A.P., L.R and S.A.S processed and analysed spatial data. M.J.R., H.A.L, and T. E. S  
420 conducted genetic analyses, and S.J.B, M.E.H and M.J.R processed and analysed  
421 isotope data. M.J.R wrote paper and all authors discussed the results and contributed to  
422 the manuscript.

423

424 **Competing financial interests:** The authors declare no competing financial interests.

425

426

427

428

429

430

431

432

433

434

435

436 **Figure legends**

437 **Fig. 1. Pacific migrations of tracked *P. cookii***

438 Movements of *P. cookii* breeding on Little Barrier Island (red circle, North Island) and  
439 Codfish Island (purple circle, South Island), New Zealand. Approximate post-breeding  
440 and pre-breeding migration routes of tracked LBI (red dashed lines) and CDF (purple  
441 dashed lines) birds begin and end with mean migration departure and arrival dates  
442 (white text). Red and purple tones, and associated contour lines, represent the 95%,  
443 75%, 50%, and 25% kernel estimates for LBI and CDF birds respectively. Pie charts  
444 show the geographic distribution of cytochrome *c* oxidase subunit 1 mtDNA haplotypes  
445 sequenced from modern *P. cookii* populations (charts attached to LBI and CDF colony  
446 locations) and historical *P. cookii* skins collected in the North (BC) and South Pacific  
447 (HC) Ocean in 1905 and 1913 respectively (charts attached to white circles showing  
448 approximate collection location). Pie chart size reflects genetic analysis sample size:  
449 modern *P. cookii* LBI n = 26 and CDF n = 19; historical *P. cookii* BC n = 9 and HC n =  
450 10.

451

452 **Fig. 2 Isotope ratios of contemporary and historic *P. cookii* feathers**

453 Feather isotope signatures of *P. cookii* ( $\pm$ SD) from LBI (red circle, 2006 (untracked)  
454 and 2008 (tracked) combined, n = 31) and CDF (purple circle, 2006 (untracked) and

455 2008 (tracked) combined,  $n = 29$ ) and from historical skins collected in the North  
456 Pacific (BC) (black circle, 1905,  $n = 9$ ) and South Pacific Ocean (HC) (clear circle,  
457 1913,  $n = 10$ ).

458

459 **Fig. 3 Characteristics of selected oceanic habitats**

460 Box plots of median values for remotely sensed environmental data (a. water depth, b.  
461 sea surface temperature, c. chlorophyll *a* concentration) from logger locations of *P.*  
462 *cooki* tracked from Little Barrier Island (LBI,  $n = 11$ ) and Codfish Island (CDF,  $n = 11$ )  
463 during occupancy of core non-breeding habitats (LBI Pres. and CDF Pres., 20% UD)  
464 and in randomly selected proximate locations (LBI Abs. and CDF Abs.) see Methods.  
465 Asterisks represent significance of difference for group comparisons: \*  $p = 0.05$ , \*\*  $p =$   
466  $0.01$ , \*\*\*  $p = 0.001$ , - = not significant. Box plots illustrate 25<sup>th</sup>, 50<sup>th</sup> (median), and  
467 75th percentiles, error bars represent minimum and maximum values falling within 1.5  
468 Inter Quartile Range, and crosses plot the outliers.

469

470

471

472 **References**

- 473 1 Dor, R., Safran, R. J., Sheldon, F. H., Winkler, D. W. & Lovette, I. J. Phylogeny  
 474 of the genus *Hirundo* and the barn swallow subspecies complex. *Molecular*  
 475 *Phylogenetics and Evolution* **56**, 409-418, doi:DOI:  
 476 10.1016/j.ympev.2010.02.008 (2010).
- 477 2 Irwin, D. E. & Irwin, J. H. in *Birds of Two Worlds: The Ecology and Evolution*  
 478 *of Migration* (eds R Greenberg & P P Marra) 27-40 (Johns Hopkins University  
 479 Press, Baltimore, 2005).
- 480 3 Mayr, E. *Systematics and the Origin of Species*. (Columbia University Press,  
 481 1942).
- 482 4 Irwin, D. E., Bensch, S., Irwin, J. H. & Price, T. D. Speciation by distance in a  
 483 ring species. *Science* **307**, 414-416, doi:10.1126/science.1105201 (2005).
- 484 5 Bearhop, S. *et al.* Assortative mating as a mechanism for rapid evolution of a  
 485 migratory divide. *Science* **310**, 502-504 (2005).
- 486 6 Sorenson, M. D., Sefc, K. M. & Payne, R. B. Speciation by host switch in brood  
 487 parasitic indigobirds. *Nature* **424**, 928-931,  
 488 doi:[http://www.nature.com/nature/journal/v424/n6951/supinfo/nature01863\\_S1](http://www.nature.com/nature/journal/v424/n6951/supinfo/nature01863_S1.html)  
 489 [html](http://www.nature.com/nature/journal/v424/n6951/supinfo/nature01863_S1.html) (2003).
- 490 7 Galarza, J. A. *et al.* The influence of oceanographic fronts and early-life-history  
 491 traits on connectivity among littoral fish species. *Proceedings of the National*  
 492 *Academy of Sciences* **106**, 1473-1478, doi:10.1073/pnas.0806804106 (2009).
- 493 8 Friesen, V. L., Burg, T. M. & McCoy, K. D. Mechanisms of population  
 494 differentiation in seabirds. *Molecular Ecology* **16**, 1765-1785,  
 495 doi:10.1111/j.1365-294X.2006.03197.x (2007).
- 496 9 Natoli, A., Birkun, A., Aguilar, A., Lopez, A. & Hoelzel, A. R. Habitat structure  
 497 and the dispersal of male and female bottlenose dolphins (*Tursiops truncatus*).  
 498 *Proceedings of the Royal Society of London B: Biological Sciences* **272**, 1217-  
 499 1226, doi:10.1098/rspb.2005.3076 (2005).
- 500 10 Egevang, C. *et al.* Tracking of Arctic terns *Sterna paradisaea* reveals longest  
 501 animal migration. *Proceedings of the National Academy of Sciences of the*  
 502 *United States of America* **107**, 2078-2081, doi:10.1073/pnas.0909493107  
 503 (2010).
- 504 11 Shaffer, S. A. *et al.* Migratory shearwaters integrate oceanic resources across the  
 505 Pacific Ocean in an endless summer. *Proceedings of the National Academy of*  
 506 *Sciences of the United States of America* **103**, 12799-12802,  
 507 doi:10.1073/pnas.0603715103 (2006).
- 508 12 Burg, T. M. & Croxall, J. P. Global relationships amongst black-browed and  
 509 grey-headed albatrosses: analysis of population structure using mitochondrial  
 510 DNA and microsatellites. *Molecular Ecology* **10**, 2647-2660 (2001).
- 511 13 Austin, J., White, R. & Ovenden, J. Population-genetic structure of a philopatric,  
 512 colonial nesting seabird, the short-tailed shearwater (*Puffinus tenuirostris*) *Auk*  
 513 **111**, 70-79 (1994).
- 514 14 Avise, J. C., Nelson, W. S., Bowen, B. W. & Walker, D. Phylogeography of  
 515 colonially nesting seabirds, with special reference to global matrilineal patterns  
 516 in the sooty tern (*Sterna fuscata*) *Molecular Ecology* **9** (2000).
- 517 15 Techow, N. M. S. M. *et al.* Speciation and phylogeography of giant petrels  
 518 *Macronectes*. *Molecular Phylogenetics and Evolution* **54**, 472-487, doi:DOI:  
 519 10.1016/j.ympev.2009.09.005 (2010).

- 520 16 Morris-Pocock, J. A., Steeves, T. E., Estela, F. A., Anderson, D. J. & Friesen, V.  
521 L. Comparative phylogeography of brown (*Sula leucogaster*) and red-footed  
522 boobies (*S. sula*): The influence of physical barriers and habitat preference on  
523 gene flow in pelagic seabirds. *Molecular Phylogenetics and Evolution* **54**, 883-  
524 896, doi:DOI: 10.1016/j.ympev.2009.11.013 (2010).
- 525 17 Gómez-Díaz, E., González-Solís, J. & Peinado, M. A. Population structure in a  
526 highly pelagic seabird, the Cory's shearwater *Calonectris diomedea*: an  
527 examination of genetics, morphology and ecology *Marine Ecology Progress*  
528 *Series* **382**, 197-209 (2009).
- 529 18 Dearborn, D. C., Anders, A. D., Schreiber, E. A., Adams, R. M. M. & Mueller,  
530 U. G. Inter-island movements and population differentiation in a pelagic seabird.  
531 *Molecular Ecology* **12**, 2835-2843 (2003).
- 532 19 Steeves, T. E., Anderson, D. J. & Friesen, V. L. A role for nonphysical barriers  
533 to gene flow in the diversification of a highly vagile seabird, the masked booby  
534 (*Sula dactylatra*). *Molecular Ecology* **14**, 3877-3887 (2005).
- 535 20 Burger, E. A. & Shaffer, S. A. Applications of tracking and data-logging  
536 technology in research and conservation of seabirds. *The Auk* **125**, 253-264  
537 (2008).
- 538 21 Phillips, R. A., Croxall, J. P., Silk, J. R. D. & Briggs, D. R. Foraging ecology of  
539 albatrosses and petrels from South Georgia: two decades of insights from  
540 tracking technologies. *Aquatic Conservation: Marine and Freshwater*  
541 *Ecosystems* **17**, 6-21 (2008).
- 542 22 Imber, M. J., West, J., A. & Cooper, W. J. Cook's petrel (*Pterodroma cookii*):  
543 historic distribution, breeding biology, and effects of predators. *Notornis* **50**,  
544 221-230 (2003).
- 545 23 Rayner, M. J. *et al.* Predictive habitat modelling improves the population census  
546 accuracy of a burrowing seabird: a study of the endangered Cook's petrel.  
547 *Biological Conservation* **138**, 235-247 (2007).
- 548 24 Rayner, M. J., Parker, K. A. & Imber, M. J. Population census of Cook's petrel  
549 *Pterodroma cookii* breeding on Codfish Island (New Zealand) and the global  
550 conservation status of the species. *Bird Conservation International* **18**, 211-218  
551 (2008).
- 552 25 Rayner, M. J. *et al.* Foraging ecology of the Cook's petrel *Pterodroma cookii*  
553 during the austral breeding season: a comparison of its two populations. *Marine*  
554 *Ecology Progress Series* **370**, 271-284 (2008).
- 555 26 Marchant, S. & Higgins, P. J. *Handbook of Australasian, Antarctic and New*  
556 *Zealand Birds. Volume 1 Ratites to Ducks.* (Oxford University Press, 1990).
- 557 27 Murphy, R. C. On *Pterodroma cookii* and its allies. *American Museum Novitates*  
558 **370** (1929).
- 559 28 Falla, R. A. Review of the smaller Pacific forms of *Pterodroma* and *Cookilaria*.  
560 *Emu* **42**, 111-118 (1942).
- 561 29 Gonzalez-Solis, J., Croxall, J. P., Oro, D. & Ruiz, X. Trans-equatorial migration  
562 and mixing in the wintering areas of a pelagic seabird. *Frontiers in Ecology and*  
563 *the Environment* **5**, 297-301, doi:doi:10.1890/1540-  
564 9295(2007)5[297:TMAMIT]2.0.CO;2 (2007).
- 565 30 Landers, T. J., Rayner, M. J., Phillips, R. A. & Hauber, M. E. Dynamics of  
566 seasonal movements by a trans-pacific migrant, the westland petrel. *Condor* **113**,  
567 71-79 (2011).
- 568 31 Ismar, S. M. H., Phillips, R. A., Rayner, M. J. & Hauber, M. E. Geolocation  
569 tracking of the annual migration of adult australasian gannets (*Morus serrator*)

- 570 breeding in New Zealand. *The Wilson Journal of Ornithology* **123**, 121-125  
571 (2011).
- 572 32 Rayner, M. J., Taylor, G. A., Thompson, D. R., Torres, L. G., Sagar, P. M.,  
573 Shaffer, S. A. Migration and diving activity in three post-breeding flesh-footed  
574 shearwaters (*Puffinus carneipes*). *Journal of Avian Biology* (in press).
- 575 33 Gill, R. E. *et al.* Extreme endurance flights by landbirds crossing the Pacific  
576 Ocean: ecological corridor rather than barrier? *Proceedings of the Royal Society*  
577 *B: Biological Sciences* **276**, 447-457, doi:10.1098/rspb.2008.1142 (2009).
- 578 34 Spear, L. B., Howell, S. N. G. & Ainley, D. G. Notes on the at-sea identification  
579 of some Pacific gadfly petrels (Genus: *Pterodroma*). *Colonial Waterbirds* **15**,  
580 202-218 (1992).
- 581 35 Hobson, K. A. Tracing origins and migration of wildlife using stable isotopes: a  
582 review. *Oecologia* **120**, 314-326 (1999).
- 583 36 Phillips, R., Bearhop, S., McGill, R. & Dawson, D. Stable isotopes reveal  
584 individual variation in migration strategies and habitat preferences in a suite of  
585 seabirds during the nonbreeding period. *Oecologia* **160**, 795-806,  
586 doi:10.1007/s00442-009-1342-9 (2009).
- 587 37 Cherel, Y., Phillips, R. A., Hobson, K. A. & McGill, R. Stable isotope evidence  
588 of diverse species-specific and individual wintering strategies in seabirds.  
589 *Biology Letters* **2**, 301-303, doi:10.1098/rsbl.2006.0445 (2006).
- 590 38 Rayner, M. J., Carragher, C. J. F. & Hauber, M. E. Mitochondrial DNA analysis  
591 reveals genetic structure in two New Zealand Cook's petrel (*Pterodroma cookii*)  
592 populations. *Conservation genetics* **11**, 2073-2077 (2010).
- 593 39 Monteiro, L. R. & Furness, R. W. Speciation through temporal segregation of  
594 Madeiran storm petrel (*Oceanodroma castro*) populations in the Azores?  
595 *Proceedings of the Royal Society of London B: Biological Sciences* **353**, 945-  
596 953, doi:10.1098/rstb.1998.0259 (1998).
- 597 40 Friesen, V. L. *et al.* Sympatric speciation by allochrony in a seabird.  
598 *Proceedings of the National Academy of Sciences of the United States of*  
599 *America* **104**, 18589-18594, doi:10.1073/pnas.0700446104 (2007).
- 600 41 Marra, P. P., Hobson, K. A. & Holmes, R. T. Linking winter and summer events  
601 in a migratory bird by using stable-carbon isotopes. *Science* **282**, 1884-1886,  
602 doi:10.1126/science.282.5395.1884 (1998).
- 603 42 Irwin, D. E. Speciation: new migratory direction provides route toward  
604 divergence. *Current Biology* **19**, 1111-1113, doi:DOI:  
605 10.1016/j.cub.2009.11.011 (2009).
- 606 43 Berthold, P., Helbig, A. J., Mohr, G. & Querner, U. Rapid microevolution of  
607 migratory behaviour in a wild bird species. *Nature* **360**, 668-670 (1992).
- 608 44 Phillips, R. A., Silk, J. R. D., Phalan, B., Catry, P. & Croxall, J. P. Seasonal  
609 sexual segregation in two *Thalassarche* albatrosses: competitive exclusion,  
610 reproductive role specialization or foraging niche divergence? *Proceedings of*  
611 *the Royal Society B: Biological Sciences* **271**, 1283-1292 (2004).
- 612 45 Shaffer, S. A., Weimerskirch, H. & Costa, D. P. Functional significance of  
613 sexual dimorphism in wandering albatrosses, *Diomedea exulans*. *Functional*  
614 *Ecology* **15**, 203-210 (2001).
- 615 46 Weimerskirch, H., Cherel, Y., Cuenot-Chaillet, F. & Ridoux, V. Alternative  
616 foraging strategies and resource allocation by male and female wandering  
617 albatrosses. *Ecology* **78**, 2051-2063 (1997).



- 618 47 Gómez-Díaz, E., González-Solís, J., Peinado, M. A. & Page, R. D. M.  
619 Phylogeography of the *Calonectris* shearwaters using molecular and  
620 morphometric data. *Molecular Phylogenetics and Evolution* **41**, 322-332 (2006).
- 621 48 Fridolfsson, A. K. & Ellegren, H. A simple and universal method for molecular  
622 sexing of non-ratite birds. *Journal of Avian Biology* **30**, 116-221 (1999).
- 623 49 Phillips, R. A., Silk, J. R. D., Croxall, J. P., Afanasyev, V. & Briggs, D. R.  
624 Accuracy of geolocation estimates for flying seabirds. *Marine Ecology Progress  
625 Series* **266**, 265-272 (2004).
- 626 50 Guilford, T. *et al.* Migration and stopover in a small pelagic seabird, the Manx  
627 shearwater *Puffinus puffinus*: insights from machine learning. *Proceedings of the  
628 Royal Society B: Biological Sciences* **276**, 1215-1223,  
629 doi:10.1098/rspb.2008.1577 (2009).
- 630 51 Shaffer, S. A. *et al.* Spatio-temporal habitat use by breeding sooty shearwaters  
631 *Puffinus griseus*. *Marine Ecology Progress Series* **391**, 209-220 (2009).
- 632 52 Worton, B. J. Kernel methods for estimating the utilization distribution in home-  
633 range studies. *Ecology* **70**, 164-168 (1989).
- 634 53 Sheather, S. J. & Jones, M. C. A reliable data-based bandwidth selection method  
635 for kernel density estimation. *Journal of the Royal Statistical Society B* **53**, 683-  
636 690 (1991).
- 637 54 O'Reilly, J. E. *et al.* Ocean color chlorophyll algorithms for SeaWiFS. *Journal of  
638 Geophysical Research* **103**, 24937-24953, doi:10.1029/98jc02160 (1998).
- 639 55 Powell, B. S. *et al.* 4DVAR data assimilation in the intra-americas sea with the  
640 Regional Ocean Modeling System (ROMS). *Ocean Modelling* **23**, 130-145  
641 (2008).
- 642 56 Smith, W. H. F. & Sandwell, D. T. Global sea floor topography from satellite  
643 altimetry and ship depth soundings. *Science* **277**, 1956-1962,  
644 doi:10.1126/science.277.5334.1956 (1997).
- 645 57 Quillfeldt, P., Masello, J., McGill, R., Adams, M. & Furness, R. Moving  
646 polewards in winter: a recent change in the migratory strategy of a pelagic  
647 seabird? *Frontiers in Zoology* **7**, 15 (2010).
- 648 58 Patel, S., Waugh, J., Millar, C. D. & Lambert, D. M. Conserved primers for  
649 DNA barcoding historical and modern avian samples from New Zealand and  
650 Antarctic birds. *Molecular Ecology Resources* **10**, 431-438 (2009).
- 651 59 Rozen, S. & Skaletsky, H. Primer3 on the WWW for general users and for  
652 biologist programmers. *Methods in Molecular Biology* **132**, 365-386 (2000).

653

654

655


656

657

658

659

660



Trophic level ↑

