

January 2014

Additional lichen records and minerological data from metal-contaminated sites in Maine

I. D. Medeiros

A. M. Fryday

Nishanta Rajakaruna

San Jose State University, nishanta.rajakaruna@sjsu.edu

Follow this and additional works at: http://scholarworks.sjsu.edu/biol_pub



Part of the [Botany Commons](#)

Recommended Citation

I. D. Medeiros, A. M. Fryday, and Nishanta Rajakaruna. "Additional lichen records and minerological data from metal-contaminated sites in Maine" *Rhodora* (2014): 323-347.

This Article is brought to you for free and open access by the Biological Sciences at SJSU ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SJSU ScholarWorks. For more information, please contact scholarworks@sjsu.edu.

ADDITIONAL LICHEN RECORDS AND MINERALOGICAL DATA FROM METAL-CONTAMINATED SITES IN MAINE

IAN D. MEDEIROS

College of the Atlantic, 105 Eden Street, Bar Harbor, ME 04609

ALAN M. FRYDAY

Herbarium, Department of Plant Biology, Michigan State University
612 Wilson Road, East Lansing, MI 48824

¹NISHANTA RAJAKARUNA

College of the Atlantic, 105 Eden Street, Bar Harbor, ME 04609;
Unit for Environmental Sciences and Management, North-West University,
Private Bag X6001, Potchefstroom, 2520, South Africa

¹Author for Correspondence; e-mail: nrajakaruna@coa.edu

ABSTRACT. Geochemistry and mineralogy of rocks play important roles in the occurrence of individual lichen species and assembly of lichen communities. Whereas lichens of metal-enriched settings have been a focus of study for many decades, only a few such lichen inventories exist for North America. We reexamined the lichen biota of Pine Hill, a serpentine outcrop on Little Deer Isle, Maine and Callahan Mine, a copper- and zinc-enriched Superfund site in Brooksville, Maine by conducting additional field surveys and reexamining unidentified taxa from previous collections. To better characterize the substrates upon which the lichens were found, we conducted elemental analyses via x-ray fluorescence and inductively coupled plasma-mass spectrometry on rock samples collected at Pine Hill and recorded pH, electrical conductivity, and elemental concentrations via inductively coupled plasma mass spectrometry on soil samples from Callahan Mine. The re-investigation of lichens of the two metal-enriched sites resulted in the addition of 20 taxa to Pine Hill and 10 taxa to Callahan Mine. These include *Dermatocarpon leptophyllodes*, *Placynthiella hyporhoda*, *Pyrenocarpon thelostomum*, and *Vezdaea acicularis*, all recorded for the first time from New England. In addition, we report the first documented records since the late 19th to early 20th century for New England of *Porocyphus coccodes*, *Sarcosagium campestre*, and *Steinia geophana*, and the first such record for Maine for *Coccocarpia palmicola*. *Stereocaulon condensatum* and *S. subcoralloides*, both considered as rare in New England, were also collected from Callahan Mine.

Key Words: edaphic ecology, lichen ecology, Maine lichens, metal quarries, metal-tolerance, serpentine, superfund sites

Areas of soil and exposed bedrock rich in heavy metals are geologically and ecologically distinct from surrounding areas and

may support distinct floras resulting from their disjunction (O'Dell and Rajakaruna 2011). These areas can be anthropogenic in origin, such as mines where waste accumulation on the surface has dramatically changed the landscape, or naturally occurring, such as outcrops of ultramafic rock. Decades of research have suggested that these "edaphic islands" may harbor a high proportion of endemic plants that have evolved to tolerate edaphic stresses (Kay et al. 2011; Rajakaruna and Boyd 2008). Often-quoted examples include the flora of the island of New Caledonia, which has abundant ultramafic rock outcrops and far more endemic species than many larger islands (Anacker 2011), and California, which supports a high number of species endemic to specific geologies, including ultramafic rocks such as serpentinite (Kay et al. 2011; Safford et al. 2005).

Ultramafic rocks contain at least 70% ferromagnesian silicate minerals (Kruckeberg 2002); soils weathered from such rocks tend to have ratios of calcium (Ca) to magnesium (Mg) < 1 , low levels of essential nutrients such as nitrogen (N), potassium (K), and phosphorous (P), and high levels of heavy metals such as chromium (Cr), nickel (Ni), and cobalt (Co) (Brady et al. 2005). The impact of geologically unique environments on plant diversity is especially apparent on serpentinite outcrops (hereafter referred to as serpentine), where high levels of heavy metals and low essential nutrients such as Ca limit long-term habitation to those species that possess mechanisms for edaphic tolerance (Kruckeberg 1986). Serpentine environments exert strong selective pressure on plants, and adaptations to these unique edaphic conditions drive speciation and high rates of endemism (Kay et al. 2011; Kruckeberg 1986; O'Dell and Rajakaruna 2011). Because of the distinctiveness and global rarity of many of the plants found on serpentine and their potential use for the study of plant ecology and evolution (Harrison and Rajakaruna 2011), the vascular flora of serpentine outcrops have been documented extensively in western North America, New Caledonia, Cuba, and Europe (Roberts and Proctor 1992).

Like serpentine outcrops, metal mines are also edaphically stressful environments that typically have shallow and rocky soils with low levels of plant nutrients and large concentrations of heavy metals (O'Dell and Rajakaruna 2011). Plants growing on metal-enriched mine tailings, like those on serpentine outcrops, are often physiologically distinct and reproductively isolated from their relatives growing on nearby uncontaminated soil (Antonovics

2006; O'Dell and Rajakaruna 2011), providing opportunities for the study of ecological speciation (Kay et al. 2011).

Compared to vascular plants, lichens on serpentine outcrops have received less attention (Favero-Longo et al. 2004; Rajakaruna et al. 2012) and, as with vascular plants, very little of the research has been carried out in eastern North America (Harris et al. 2007; Rajakaruna et al. 2009; Reed 1986; Sirois et al. 1988). This is somewhat surprising considering that New England has received extensive lichenological study, including investigations of sites with unique geologies (see Miller et al. 2005). Brodo (1973) remarked that since there is a close relationship between lichen and substrate, edaphically extreme substrates may have unique effects on lichens. For example, cryptogams have shown high specificity on calcareous substrates (Bates 1978). Conversely, a comprehensive review of the literature on lichens of serpentine outcrops by Favero-Longo et al. (2004) showed no evidence of serpentine endemism in lichens. Lichens are apparently less affected by ultramafic substrata than vascular plants, possibly because of differences in cell wall structure that render the low level of Ca in ultramafic rocks less detrimental to lichens than to vascular plants (O'Dell and Rajakaruna 2011). Another possible explanation is that the acids produced by many lichens can form complexes with a range of elements, thus neutralizing their toxicity (Wilson 1995). However, it is possible that the physical structure of a rock is as important to the lichen biota as its mineralogy, and that climate, elevation, history of land use, and other biotic and abiotic factors may have as much, or more, influence on the lichen biota as any chemical or physical properties of the rock itself; see Rajakaruna et al. (2012) and references therein for a full discussion of lichen-metal relations.

Despite their apparent lack of substrate specificity, lichens are able to accumulate heavy metals from metal-rich substrates (Bačkor and Loppi 2009; Lambinon et al. 1964; Maquinay et al. 1961) and unique lichen ecotypes and mixtures of calcicolous and silicolous species are often found on serpentine outcrops (Favero-Longo et al. 2004; Rajakaruna et al. 2012). The potential for unique lichen assemblages on serpentine outcrops suggests that anthropogenic, metal-enriched sites may also provide similar challenges for lichens. Purvis (1993) states that the debris left by copper mining creates habitat for a variety of lichen species, including those that are rare, a factor that should be taken into consideration when old mine sites undergo reclamation.

To date, only two published studies have documented the lichen biota of metal-enriched sites and serpentine outcrops in coastal New England. Both studies deal with sites with histories of mining or quarrying in Hancock County, Maine, U.S.A. Harris et al. (2007) catalogued the lichens of Pine Hill, an abandoned peridotite quarry on Little Deer Isle. A species list for the site was presented and compared with lists of lichen species recorded in nearby Acadia National Park (Sullivan 1996) and serpentine outcrops worldwide (Favero-Longo et al. 2004). Comparison between these lists showed that Pine Hill hosted two species previously unreported from New England, three species previously unreported from Maine, and 20 species previously unreported from serpentine outcrops worldwide. The latter category included *Lobaria*, a genus not previously reported from serpentine elsewhere. Rajakaruna et al. (2011) examined Callahan Mine, a copper- and zinc-enriched Superfund site in Brooksville. This study provided a species list for the site, including information on the general ecology of each species and notes on which species were previously known to tolerate or accumulate specific heavy metals. This study did not generate any new records for Maine or New England, but the authors did note the presence of two taxa described as regionally rare or declining, and two taxa reported as copper tolerant.

The primary objective of the current study was to conduct additional field surveys at Pine Hill and Callahan Mine to find species that might have been overlooked during the previous efforts. Additionally, unidentified lichens collected by Harris et al. (2007) and Rajakaruna et al. (2011) were identified during the current study. We also investigated the elemental chemistry of the serpentine rocks at Pine Hill and the ionic content and other chemical features of the soils at Callahan Mine to better characterize substrate tolerances for the lichens we collected. We compared the lichen biota of Pine Hill and Callahan Mine to a published species list for serpentine from Québec (Sirois et al. 1988) and to that of nearby Acadia National Park, Maine (Sullivan 1996) to determine whether any species we found were previously undocumented from serpentine substrates and other habitats in northeastern North America.

MATERIALS AND METHODS

Site descriptions. Pine Hill is a former peridotite quarry occupying about 40 acres on Little Deer Isle, Hancock County,

Maine, USA (44°17'07.3"N, 68°42'06.7"W; WGS 84). The outcrop is serpentized peridotite formed during the Jurassic period (Hooke 2003). Quarry activities from approximately 1930–1960 have reduced the outcrop to a half dome and exposed extensive vertical faces of unweathered rock with a southern exposure. The quarry floor is largely serpentized gravel overlying bedrock, with a number of large boulders near the base of the outcrop. Soils on the site range from coarse gravel on the quarry floor, to humus-rich organic debris along the three sides of the outcrop, and sandy loam at the top of the outcrop. A number of ephemeral freshwater seeps exist at the site, creating moist environments in an otherwise dry landscape. For additional information on the site see Briscoe et al. (2009), Harris et al. (2007), and Pope et al. (2010).

Callahan Mine is a former intertidal open-pit mine in Brooksville, Hancock County, Maine (44°20'N, 68°48'W; WGS 84). It has been mined intermittently since 1880, the most intensive mining taking place from 1968–1972. Callahan Mine was listed as a Superfund site in 2002 (Environmental Protection Agency 2013) due to elevated levels of heavy metals, including copper (Cu), zinc (Zn), lead (Pb), and cadmium (Cd), and organic contaminants such as PCBs (Rajakaruna et al. 2011). The site is currently under remediation. For a detailed site description see Rajakaruna et al. (2011) and Mansfield et al. (2014).

Rock elemental analysis. Elemental analysis via x-ray fluorescence (XRF) was conducted on one composite rock sample each from Pine Hill (serpentine) and nearby Settlement Quarry (granite) as a comparison. An automated sequential XRF spectrometer (Advant'XP+, Thermo ARL, Switzerland) was used, and was run at 60 keV and 60 mA with a rhodium target. Elemental concentrations were determined for major (Al–Ti) and trace (As–Zr) elements. For detailed analytical methodology see Rajakaruna et al. (2012). Inductively coupled plasma-mass spectrometry (ICP-MS) was used to analyze additional trace elements that were not determined by XRF analysis. The procedure consists of a low-dilution fusion with dilithium tetraborate (Spectromelt® A-10, EM Science, Gibbstown, NJ) followed by an open-vial mixed acid digestion. Reagents were HNO₃ 69–70% (Fisher ACS plus grade), HF 48–52% (Baker ACS reagent grade), HClO₄ 67–71% (Fisher Trace Metal Grade), and H₂O₂ (Baker ACS Reagent). The HF was further purified before use by sub-boiling distillation in a teflon still. All water used was

>18 M deionized water from a Nanopure analytical grade water system (Barnstead/Thermolyne, Dubuque, IA). Powdered rock samples were mixed with an equal amount of dilithium tetraborate flux (typically 2 g), placed in a carbon crucible and fused at 1000°C in a muffle furnace for 30 min. After cooling, the resultant fusion bead was briefly ground in a carbon-steel ring mill and a 250 mg portion was weighed into a 30 ml, screw-top Teflon PFA vial for dissolution. The acid dissolution consisted of a first evaporation with HNO₃ (2ml), HF (6 ml), and HClO₄ (2 ml) at 110°C. After evaporating to dryness, the sample was wetted and the sides of the vial were rinsed with a small amount of water before a second evaporation with HClO₄ (2 ml) at 160°C. After the second evaporation, samples were brought into solution by adding approximately 10 ml of water, 3 ml HNO₃, 5 drops H₂O₂, and 2 drops HF, and warmed on a hot plate until a clear solution was obtained. The sample was then transferred to a clean 60 ml HDPE bottle and diluted up to a final weight of 60 g with deionized water. Solutions were analyzed on an Agilent Model 4500 ICP-MS and were diluted an additional 10× at the time of analysis using Agilent's Integrated Sample Introduction System (ISIS; Agilent Technologies, Santa Clara, CA). All analyses were performed at the Hooper GeoAnalytical Laboratory, Washington State University, WA, USA.

Soil chemical analysis. The lichens at Callahan Mine were collected from within or close to the plots established by Mansfield et al. (2014). Therefore, we used the same soil chemical data reported in their Table 4 (for the tailings pond, waste rock piles, and habitats 'in between') to better characterize the range of tolerance to various soil chemical features for the lichens we collected. Methods of soil analyses are reported in Mansfield et al. (2014).

Lichen identification. Lichens for this study were collected by R. C. Harris (New York Botanical Garden, NY, USA) in 2009 and by one of us (A.M.F.) in 2013. Several unidentified *Cladonia* and *Lepraria* specimens, collected in 2004 by T. B. Harris as part of the original Pine Hill lichen survey (Harris et al. 2007) and deposited at HCOA, were also identified for this project by one of us (I.D.M.). Lichens were identified by A.M.F. and I.D.M. using a variety of published keys, including Brodo et al. (2001), Gowan and Brodo (1988), and Hinds and Hinds (2007). Two collections that we were unable to identify to species (*Aspicilia* and *Lichinaceae*) were

sent to experts in these groups. Thin-layer chromatography was performed and secondary metabolites identified according to procedures and data in Orange et al. (2001). Nomenclature follows Esslinger (2012). Voucher specimens for this study and Harris et al. (2007) have been deposited at HCOA, MSC, NBM, and NY.

Lichen distribution data for North America are mostly scattered within the literature and difficult to obtain. The recent innovation of the Consortium of North American Lichen Herbaria (CNALH), which aims to digitize the label data from all lichen collections in North American herbaria, was initiated to correct this deficit and we have based our assumptions of species' distributions on records contained in this database. This is an unsatisfactory solution, but the best that is currently available. Additionally, the distribution of New England macrolichens was based on Hinds and Hinds (2007).

RESULTS

Recent collections and re-examination of previously collected lichen material have resulted in the addition of 20 taxa to the species list for Pine Hill, and 10 taxa to that of Callahan Mine (Appendix). There are now 82 lichen taxa from 40 genera known from Pine Hill and 84 lichen taxa from 41 genera known from Callahan Mine. Fifty-nine percent (48/82) of the Pine Hill taxa and 64% (54/84) of the Callahan Mine taxa are shared with the lichen biota of Acadia National Park as documented by Sullivan (1996), the most recent comprehensive lichenological survey of the Park. Fifty-two percent of Pine Hill taxa (43/82) are shared with Sirois et al. (1988), the only published study of lichens of serpentine rocks in northeastern North America besides the current study and Harris et al. (2007).

The new collections include *Dermatocarpon leptophyllodes*, *Placynthiella hyporhoda*, *Pyrenocarpon thelostomum*, and *Vezdaea acicularis*, which are all recorded for the first time from New England. In addition, since the late 19th to early 20th century for New England, this survey has furnished the first modern documented records of *Porocyphus coccodes*, *Sarcosagium campetrite*, and *Steinia geophana*, and the first such record from Maine for *Coccocarpia palmicola*.

No macrolichens collected for this study are species reported by Hinds and Hinds (2007) as globally rare. However, 10 taxa were found that, according to these authors, are rare in New England: seven from Pine Hill (*Cladonia symphycarpa*, *Coccocarpia*

palmicola, *Fuscopannaria praetermissa*, *Nephroma parile*, *Pannaria rubiginosa*, *Psorula rufonigra*, and *Spilonema revertens*) and three from Callahan Mine (*Stereocaulon condensatum*, *S. subcoralloides*, and *Leptogium imbricatum*). A further three taxa from Pine Hill, *Anaptychia palmulata*, *Cladonia boryi*, and *Parmotrema crinitum*, are uncommon and potentially declining in New England (Hinds and Hinds 2007).

Eight additional taxa collected from Pine Hill were found by Sullivan (1996) in three or fewer study plots out of the 83 locations he surveyed: *Anaptychia palmulata*, *Candelariella aurella*, *Lepraria neglecta* s. lat. (syn. *L. caesioalba*), *Pannaria rubiginosa*, *Placynthiella icmalea*, *Scoliciosporum umbrinum*, *Xanthoparmelia plittii*, and *Xanthoria elegans*. Fourteen taxa collected at Callahan Mine fulfilled this criterion for local rarity: *Aspicilia* aff. *verrucigera*, *Caloplaca microthallina*, *Candelariella aurella*, *Cladonia phyllophora*, *C. stygia*, *Montanelia soredata*, *Physcia aipolia*, *P. dubia*, *P. subtilis*, *Placynthiella icmalea*, *Rhizocarpon grande*, *Scoliciosporum umbrinum*, *Xanthoparmelia plittii*, and *Xanthoria elegans*.

Elemental analyses provided substrate chemistry data for the rocks (Table 1) and soils (Table 2) from which the lichens were collected. The rocks from Pine Hill were distinct from the more regionally abundant granite in their high concentrations of heavy metals, particularly Cr and Ni, and low concentrations of Ca compared to Mg. The granitic rocks from adjacent Settlement Quarry, which were analyzed as a comparison, contained relatively higher Al and Si and radioactive trace elements such as barium (Ba), rubidium (Rb), and strontium (Sr). The mine soils from which the lichens were collected were high in Cu, Zn, and Pb. Distinct habitats within the mine (tailings pond, waste rock piles, and 'in between') were found to differ with respect to some soil chemical features (Mansfield et al. 2014), notably higher pH, Ca, and Zn in the tailings pond, higher sulfur (S) in the waste rock piles, and higher Pb in the in between habitats.

DISCUSSION

Of the species newly reported from Pine Hill, 14 were not reported from Mont Albert by Sirois et al. (1988) and should be included in the list of species reported from serpentine from northeastern North America (Rajakaruna et al. 2009). These are *Buellia lepidastra*, *Cladonia cryptochlorophaea*, *C. dimorphoclada*,

Table 1. Elemental chemistry of serpentinite and granite rock samples collected from Pine Hill and Settlement Quarry, respectively. Presence of major elements (Al–Ti) are reported as % weight, whereas the minor elements (As–Zr) are reported as ppm. Major elements and some trace elements were determined via x-ray fluorescence analysis (XRF). Additional trace elements (indicated by *) were determined by inductively coupled plasma-mass spectrometry (ICP-MS). LOI = % loss on ignition.

Element	Concentration of Elements	
	Serpentinite	Granite
Major Elements (% weight)		
Al	6.35	14.17
Ca	3.71	2.00
Fe	9.25	2.82
K	0.06	4.43
Mg	30.25	0.92
M	0.155	0.066
Na	0.04	3.66
P	0.048	0.171
Si	38.08	70.74
Ti	88.39	99.50
Sum	0.453	0.525
LOI (%)	10.43	0.58
Trace Elements (ppm)		
Ba	30	599
Ce	4	106
Cr	2824	15
Cu	83	5
Cs*	0.58	1.94
Dy*	1.58	6.64
Er*	0.88	3.73
Eu*	0.41	1.46
Ga	7	18
Gd*	1.34	6.65
Hf*	0.81	5.45
Ho*	0.32	1.32
La*	0.82	48.30
Lu*	0.13	0.56
Nb	1.8	23.4
Nd	3	44
Ni	1527	10
Pb	0	20
Pr*	0.59	11.90
Rb	2	152
Sc	17	7
Sm*	1.05	7.38
Sr	30	240

Table 1. Continued.

Element	Concentration of Elements	
	Serpentinite	Granite
Ta*	0.02	2.20
Tb*	0.25	1.10
Th	2	22
Tm*	0.12	0.57
U*	0.02	3.60
V	85	40
Y	10	37
Yb*	0.78	3.61
Zn	67	45
Zr	33	281

Table 2. Results from soil analyses at Callahan Mine shown as the mean (\pm SE) for each soil feature. Locations: TP = tailings pond, WR = waste rock piles, IB = in between, including shore of Goose Pond; N = number of samples per location. Elemental concentrations: EC is reported as mmhos/cm and all other elemental concentrations are reported as mg/kg (ppm). LOI = loss on ignition (%).

Soil Feature	Elemental Concentrations at 3 Locations		
	TP (N = 3)	WR (N = 6)	IB (N = 5)
pH	7.5 (\pm 0.1)	5.1 (\pm 0.6)	5.9 (\pm 0.1)
% LOI	0.6 (\pm 0.2)	1.1 (\pm 0.1)	5.2 (\pm 1.8)
EC	0.8 (\pm 0.3)	0.7 (\pm 0.1)	0.7 (\pm 0.2)
NO ₃ -N	1.8 (\pm 0.4)	1.4 (\pm 0.3)	6.2 (\pm 3.1)
NH ₄ -N	3.7 (\pm 0.6)	2.0 (\pm 0.1)	8.2 (\pm 5.9)
Ca	16285.0 (\pm 1514.9)	2411.5 (\pm 1291.7)	2568.6 (\pm 1429.0)
K	11.7 (\pm 0.7)	10.5 (\pm 1.4)	105.7 (\pm 41.7)
Mg	100.6 (\pm 10.7)	66.7 (\pm 9.4)	102.4 (\pm 32.3)
Na	5.6 (\pm 1.5)	5.8 (\pm 0.7)	18.4 (\pm 6.3)
P	20.1 (\pm 1.8)	4.1 (\pm 1.7)	7.3 (\pm 1.2)
S	89.8 (\pm 29.2)	376.7 (\pm 109.8)	164.1 (\pm 73.2)
Al	0.4 (\pm 0.0)	9.5 (\pm 5.8)	12.4 (\pm 11.6)
Cd	3.0 (\pm 0.9)	1.3 (\pm 0.5)	2.7 (\pm 0.7)
Cr	0.0 (\pm 0.0)	0.0 (\pm 0.0)	0.1 (\pm 0.0)
Cu	102.6 (\pm 7.2)	107.7 (\pm 25.8)	131.6 (\pm 26.4)
Fe	2.5 (\pm 0.9)	35.8 (\pm 15.6)	117.2 (\pm 62.9)
Mn	0.8 (\pm 0.2)	1.8 (\pm 0.7)	4.2 (\pm 2.0)
Mo	0.5 (\pm 0.0)	0.3 (\pm 0.1)	0.4 (\pm 0.1)
Ni	0.5 (\pm 0.2)	0.5 (\pm 0.1)	1.5 (\pm 0.6)
Pb	9.2 (\pm 2.7)	13.2 (\pm 5.7)	129.1 (\pm 78.1)
Zn	749.8 (\pm 51.9)	368.5 (\pm 118.6)	558.5 (\pm 110.2)

C. grayi, *Coccocarpia palmicola*, *Dermatocarpon leptophyllodes*, *Lecidella patavina*, *Lepraria finkii*, *Physcia tenella*, *Placidium squamulosum*, *Porocyphus coccodes*, *Pyrenocarpon thelostomum*, *Rhizocarpon disparum*, and *Xanthoparmelia viriduloumbrina*.

Of the lichen taxa newly reported from Pine Hill, *Coccocarpia palmicola* is a mainly tropical species that is known from isolated localities along the east coast of North America as far north as Newfoundland (CNALH 2013). Our collection is only the second record of *C. palmicola* in New England since the late 19th to early 20th century, and only the second ever in Maine (Hinds and Hinds 2007). This species has declined in its New England distribution over the past century (Hinds and Hinds 2007), possibly due to the susceptibility of cyanolichens to increased air pollution (Richardson and Cameron 2004). *Nephroma parile* and *Pannaria rubiginosa*, two regionally rare cyanolichens (Hinds and Hinds 2007) from Pine Hill, may be in decline for the same reason.

Dermatocarpon leptophyllodes is possibly new to eastern North America because it was not included in Hinds and Hinds (2007) and all the records in CNALH (2013) are from western states, whereas *Pyrenocarpon thelostomum* has previously been reliably reported from North America only from Illinois (M. Schultz, Biozentrum Klein Flottbek und Botanischer Garten der Universität Hamburg, Hamburg, Germany, pers. comm.), not Michigan as reported by Dillman et al. (2012). Both were found at Pine Hill in a calcareous seep along with *Porocyphus coccodes*, which is apparently the first recent record for New England since it was collected by H. Willey in Massachusetts in the 19th century (CNALH 2013). Similarly, *Coccocarpia palmicola*, which has been found on calcareous substrates in Massachusetts (Hinds and Hinds 2007), and other species known to favor substrates with a high pH, was also found at Pine Hill. This is presumably the result of preferential leaching of basic cations from the serpentine substrate, a similar effect to that reported by Miller et al. (2005) from granitic rocks on Katahdin. *Coccocarpia palmicola*, however, is usually a corticolous species and its occurrence in a basic seep on serpentine is most probably a result of the additional nutrients provided by this habitat that allow it to occur outside its normal range. Two other Pine Hill lichens (*Cladonia symphycarpa* and *Fuscopannaria praetermissa*), considered as rare in New England (Hinds and Hinds 2007), are also calcicoles.

Rarity is not a trait exclusive to calcicoles at Pine Hill. *Psorula rufonigra* and *Spilonema revertens* are two intimately linked lichen taxa, with *P. rufonigra* occurring only on the thallus of *S. revertens*. They usually occur on acidic rocks but have previously been documented in northern New England only on serpentine in Vermont (Hinds and Hinds 2007), and were again reported from serpentine at Pine Hill. Other uncommon calcifuge species include *Anaptychia palmulata* and *Parmotrema crinitum*, which are described in Hinds and Hinds (2007) as uncommon, and *Cladonia boryi*, which has experienced a range contraction in New England over the past century; it is common on the coast of Maine, but uncommon elsewhere (Hinds and Hinds 2007).

Of the species newly reported from Callahan Mine, *Placynthiella hyporhoda*, *Sarcosagium campestre*, *Steinia geophana*, and *Vezdaea acicularis* are inconspicuous terricolous species that are almost certainly under recorded. However, only *Sarcosagium campestre* and *Steinia geophana* have previously been documented from New England: *S. campestre* from a 1938 record from Vermont and *S. geophana* from a 19th century record from Massachusetts (CNALH 2013). The two, newly reported *Stereocaulon* species (*S. condensatum* and *S. subcoralloides*) are also both terricolous species, and although both have previously been reported from Maine, both are rare with only scattered occurrences (Hinds and Hinds 2007). Although not restricted to soils with a high mineral content, all these species are often found associated with disused heavy metal mines (Plantlife 2012; Purvis and Halls 1996) and the occurrence of these rare terricolous species at Callahan Mine is probably an indirect effect of the high metal content of the soil that suppresses vascular plant growth and provides a suitable substrate for their development.

Two collections from the waste rock pile at Callahan Mine were clearly referable to the genus *Amandinea* (pigmented 1-septate ascospores, long curved conidia) but could not confidently be identified to species. The immature ascospores had the thickened septum typical of *A. pelidna* but the thallus was not as well developed as is usual for this species and the ascospores were also smaller than is typical for both *A. pelidna* and *A. punctata*. They are treated here, along with previous collections, as *A. punctata* aggr.

Also collected from Callahan Mine was an *Aspicilia* sp. similar to *A. verrucigera* but with smaller ascospores [$13\text{--}15(-16) \times 8\text{--}10 \mu\text{m}$]

and conidia [(11–)12–13(–15) μm long], both with a limited range, that is probably con-specific with '*A. verrucigeroides*' (I. Brodo, Canadian Museum of Nature, pers. comm.). Molecular analysis will be undertaken to determine the systematic position of this collection. As *A. verrucigera* has previously been reported from this site we have not included this as a new taxon, but changed the listing to *A. aff. verrucigera* in Appendix.

Leptogium imbricatum is a rare calcicolous species that was collected at Callahan Mine near Goose Pond. Although we do not have a pH value for the exact location of this collection, a pH of 5.9 was reported for the 'in-between' soils (Table 2), which included soils collected along Goose Pond. Rajakaruna et al. (2011) attribute the presence of this and other calcicolous species at Callahan Mine to mortar and concrete debris left by the mining operations. Also collected from near Goose Pond was *Physcia subtilis*, which is a species of non-calcareous rocks, often in sites with sunlight exposure (Brodo et al. 2001). Maine is at the northernmost edge of its range and it was represented in Sullivan (1996) by a single specimen found on an acidic cliff face. The neutral to slightly basic pH at the tailings pond, as compared to the low pH values recorded for the waste rock piles and 'in-between' area, does not exert a discernible effect on the distribution of lichens at Callahan Mine. However, the four taxa reported from only the tailings pond, *Peltigera canina*, *P. rufescens*, *Sarcosagium campestre*, and *Steinina geophana*, all occur primarily on bare, damp ground and are typical of recently disturbed soils, although only *P. rufescens* is reported to occur mainly on basic substrata. Although it is difficult to attribute any one edaphic factor to the presence of a lichen species, it is likely that the fine-textured (silt-clay), metal-enriched, and water-logged soils at the tailings pond have created a unique habitat for some soil-crust colonizing lichens.

Five of the species newly reported from Callahan Mine (all from damp soil)—*Placynthiella hyperhoda*, *Sarcosagium campestre*, *Steinera geophana*, *Stereocaulon condensatum*, and *Vezdaea acicularis*—are considered indicators of an important site for metal-tolerant lichens (Plantlife 2012). With the addition of *Acarospora sinopica*, *Stereocaulon dactylophyllum*, and *S. pileatum*, which were already reported from the site, there are now eight taxa considered indicators. Although the Plantlife indicator list was intended for metalliferous sites in Wales, it is still relevant to eastern North America and, in fact, a further species, *Stereocaulon*

subcoralloides, should probably be added to the list for eastern North America. The presence of three indicator species is considered to be sufficient to designate a site as important for metal-tolerant lichens (Plantlife (2012), so the occurrence of eight indicator species at Callahan Mine is indicative of a very important site.

Although the importance of rock mineralogy, including elemental geochemistry, in determining the composition of saxicolous lichen communities has long been recognized (Purvis and Halls 1996), attempts to analyze the distribution of saxicolous lichens according to their lithochemistry are not very common (Brodo 1973; Werner 1956). Further, studies that directly associate mineralogy or elemental chemistry of host rocks to the presence of lichen species or the assemblage of lichen communities are extremely rare (Boyle et al. 1987; Rajakaruna et al. 2012). Although complex interactions between lichens and rocks and lichens and elements are often noted (Hauck et al. 2007; Purvis 1996; Richardson 1995; Shimizu 2004; Wilson 1995), the exact nature of this substratum-level influence on lichens (i.e., whether chemical and/or textural) is unclear (Rajakaruna et al. 2012). Our sampling strategy does not allow us to directly associate the presence of any lichen species that we collected with the concentration of a particular element or combination of elements. However, our analyses of substrate chemistry of the rocks and soils from which these lichens were collected have improved the characterization of the range of elemental tolerances for the lichens found at Pine Hill and Callahan Mine.

Our study has highlighted the importance of metal rich rocks and soils in harboring rare and unusual lichen species and assemblages. An informal survey by the authors of the granitic Settlement Quarry in the vicinity of Pine Hill suggests that there are major differences between the two sites with regard to the most prevalent saxicolous species (Medeiros et al., unpubl. data). However, it is unclear whether these differences result from the chemical differences we have reported (Table 1) or from surface textural differences observed between granitic and serpentinite rocks. Surveys of geographically and climatically similar areas with different geologies could help answer the question of whether the number of regionally rare or uncommon lichen species at Pine Hill and Callahan Mine can be attributed to chemical or physical features of the substrate, or whether other factors not directly

related to geology are the dominant forces at work. Such surveys could also reaffirm the importance of conserving diverse habitats to protect the diversity of lichens. Additional work to document lichens on serpentine outcrops and mine sites elsewhere that are exposed to a wide range of climatic and edaphic variation will help document the lichen biota of understudied habitats and contribute to our understanding of the effects of heavy metals and harsh environments on lichen diversity.

ACKNOWLEDGMENTS. The authors thank the Island Heritage Trust of Deer Isle for permission to collect at Pine Hill and for continued support of our research activities at the Pine Hill Preserve and Settlement Quarry; Naji Akladiss (U.S. Environmental Protection Agency) and Sally N. Mills (Hale & Hamlin, LLC) for providing access to Callahan Mine; Richard Harris (New York Botanical Garden) for collections made at Pine Hill; Irwin M. Brodo (Canadian Museum of Nature, Ottawa) and James C. Lendemer (New York Botanical Garden) for help with a difficult *Lepraria* specimen; Björn Owe-Larsson (Herbarium, Botany Section, Museum of Evolution, Uppsala, Sweden) and Matthias Schultz (Biozentrum Klein Flottbek und Botanischer Garten der Universität Hamburg, Hamburg, Germany) for identifying specimens of *Aspicilia* and Lichinaceae (*Porocyphus* and *Pyrenocarpon*), respectively; and Irwin M. Brodo, and two anonymous reviewers for their careful reviews that greatly improved the paper.

LITERATURE CITED

- ANACKER, B. L. 2011. Phylogenetic patterns of endemism and diversity, pp. 49–79. *In*: S. P. Harrison and N. Rajakaruna, eds., *Serpentine: The Evolution and Ecology of a Model System*. University of California Press, Berkeley, CA.
- ANTONOVICS, J. 2006. Evolution in closely adjacent plant populations. X. Long-term persistence of prereproductive isolation at a mine boundary. *Heredity* 97: 33–37.
- BAČKOR, M. AND S. LOPPI. 2009. Interactions of lichens with heavy metals. *Biol. Pl.* 53: 214–222.
- BATES, J. W. 1978. The influence of metal availability on the bryophyte and macrolichen vegetation of four rock types on Skye and Rhum. *J. Ecol.* 66: 457–482.
- BOYLE, A. P., P. M. MCCARTHY, AND D. STEWART. 1987. Geochemical control of saxicolous lichen communities on the Creggaun Gabbro, Letterfrack, Co. Galway, western Ireland. *Lichenologist* 19: 307–317.

- BRADY, K. U., A. R. KRUCKEBERG, AND H. D. BRADSHAW JR. 2005. Evolutionary ecology of plant adaptation to serpentine soils. *Annual Rev. Ecol. Evol. Syst.* 36: 243–266.
- BRISCOE, L. R. E., T. B. HARRIS, E. DANNENBERG, W. BROUSSARD, F. C. OLDAY, AND N. RAJAKARUNA. 2009. Bryophytes of adjacent serpentine and granite outcrops on the Deer Isles, Maine, USA. *Rhodora* 111: 1–20.
- BRODO, I. M. 1973. Substrate ecology, pp. 410–441. *In*: V. Ahmadjian and M. E. Hale, eds., *The Lichens*. Academic Press, New York, NY.
- , S. D. SHARNOFF, AND S. SHARNOFF. 2001. *Lichens of North America*. Yale University Press, New Haven, CT.
- CNALH. 2013. Consortium of North American Lichen Herbaria. CNALH Administration at Arizona State University, Tempe, AZ, Website (<http://lichenportal.org/portal/index.php>). Accessed 5 Dec 2013.
- DILLMAN, K. L., T. AHTI, AND C. R. BJÖRK, et al. (2012). New records, range extensions, and nomenclatural innovations for lichens and lichenicolous fungi from Alaska, U.S.A. *Herzogia* 25: 177–210.
- ENVIRONMENTAL PROTECTION AGENCY. 2013. Waste site cleanup and reuse in New England. U.S. EPA, Region 1, Boston, MA. Website (<http://www.epa.gov/region1/superfund/sites/callahan>). Accessed 8 Dec 2013.
- ESSLINGER, T. L. 2012. A cumulative checklist for the lichen-forming, lichenicolous, and allied fungi of the continental United States and Canada. North Dakota State University, Fargo, ND, Website (<http://www.ndsu.nodak.edu/instruct/esslinge/chcklst/chcklst7.htm>). Accessed 26 Oct 2013.
- FAVERO-LONGO, S. E., D. ISOCRONSO, AND R. PIERVITTORI. 2004. Lichens and ultramafic rocks: A review. *Lichenologist* 36: 391–404.
- GOWAN, S. P. AND I. M. BRODO. 1988. The lichens of Fundy National Park, New Brunswick, Canada. *Bryologist* 91: 255–325.
- HARRIS, T. B., F. C. OLDAY, AND N. RAJAKARUNA. 2007. Lichens of Pine Hill, a peridotite outcrop in eastern North America. *Rhodora* 109: 430–447.
- HARRISON, S. P. AND N. RAJAKARUNA, eds. 2011. *Serpentine: The Evolution and Ecology of a Model System*. University of California Press, Berkeley, CA.
- HAUCK, M., S. HUNECK, J. A. ELIX, AND A. PAUL. 2007. Does secondary chemistry enable lichens to grow on iron-rich substrates? *Flora* 202: 471–478.
- HINDS, J. W. AND P. L. HINDS. 2007. *The Macrolichens of New England*. New York Botanical Garden Press, New York, NY.
- HOOKE, R. LEB. 2003. *A Geologic History of Deer Isle, Maine: A Thumb Nail Sketch*. Published by the author, Orono, ME.
- INDEX FUNGORUM PARTNERSHIP. 2014. *Index Fungorum*. A community resource. CABI, CBS, and Landcare Research, custodians. CABI, Wallingford, Oxfordshire, U.K.; CBS KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands; and Manaaki Whenua - Landcare Research, Lincoln, New Zealand. Website (<http://www.indexfungorum.org>). Accessed Jun 2014.
- KAY, K. M., K. L. WARD, L. R. WATT, AND D. W. SCHEMSKE. 2011. Plant speciation, pp. 97–137. *In*: S. P. Harrison and N. Rajakaruna, eds., *Serpentine: Evolution and Ecology in a Model System*. University of California Press, Berkeley, CA.

- KRUCKEBERG, A. R. 1986. An essay: The stimulus of unusual geologies for plant speciation. *Syst. Bot.* 11: 455–463.
- . 2002. *Geology and Plant Life: The Effects of Landforms and Rock Type on Plants*. University of Washington Press, Seattle, WA.
- LAMBINON, J., A. MAQUINAY, AND J. L. RAMAUT. 1964. La teneur en zinc de quelques lichens des terrains calaminaires Belges. *Bull. Jard. Bot. de l'État. (Bruxelles)* 34: 273–282.
- MANSFIELD, M., N. POPE, G. MITTLEHAUSER, AND N. RAJAKARUNA. 2014. Diversity and soil-tissue elemental relations of vascular plants of Callahan Mine, Brooksville, Maine, USA. *Rhodora* 116: xxx–yyy.
- MAQUINAY, A., I. M. LAMB, J. LAMBINON, AND J. L. RAMAUT. 1961. Dosage du zinc chez un lichen calaminaire Belge: *Stereocaulon nanodes* Tuck. f. *tyroliense* (Nyl.) M. Lamb. *Physiol. Pl. (Copenhagen)* 14: 284–289.
- MILLER, N. G., A. M. FRYDAY, AND J. W. HINDS. 2005. Bryophytes and lichens of a calcium-rich spring seep isolated on the granitic terrain of Mt. Katahdin, Maine, U.S.A. *Rhodora* 107: 339–358.
- O'DELL, R. E. AND N. RAJAKARUNA. 2011. Intraspecific variation, adaptation, and evolution, pp. 97–137. *In*: S. P. Harrison and N. Rajakaruna, eds., *Serpentine: Evolution and Ecology in a Model System*. University of California Press, Berkeley, CA.
- ORANGE, A., P. W. JAMES, AND F. J. WHITE. 2001. *Microchemical Methods for the Identification of Lichens*. British Lichen Society, London, UK.
- PLANTLIFE. 2012. *Heavy-metal Lichens in Wales. A Management Guide*. Plantlife Cymru, Bangor, Gwynedd, Wales, UK.
- POPE, N., T. B. HARRIS, AND N. RAJAKARUNA. 2010. Vascular plants of adjacent serpentine and granite outcrops on the Deer Isles, Maine, U.S.A. *Rhodora* 112: 105–141.
- PURVIS, O. W. 1993. The botanical interest of mine spoil heaps: The lichen story. *J. Russell Soc.* 5: 45–48.
- . 1996. Interactions of lichens with metals. *Sci. Progr.* 79: 283–309.
- AND C. HALLS. 1996. A review of lichens in metal-enriched environments. *Lichenologist* 28: 571–601.
- RAJAKARUNA, N. AND R. S. BOYD. 2008. The edaphic factor, pp. 1201–1207. *In*: S. E. Jorgensen and B. Fath, eds., *The Encyclopedia of Ecology*, Vol. 2. Elsevier, Oxford, U.K.
- , T. B. HARRIS, AND E. B. ALEXANDER. 2009. Serpentine geoecology of eastern North America: A review. *Rhodora* 111: 21–108.
- , ———, S. R. CLAYDEN, A. C. DIBBLE, AND F. C. OLDA. 2011. Lichens of the Callahan Mine, a copper- and zinc-enriched superfund site in Brooksville, Maine, U.S.A. *Rhodora* 113: 1–31.
- , K. KNUDSEN, A. M. FRYDAY, R. E. O'DELL, N. POPE, F. C. OLDA, AND S. WOOLHOUSE. 2012. Investigation of the importance of rock chemistry for saxicolous lichen communities of the New Idria serpentinite mass, San Benito County, California, USA. *Lichenologist* 44: 695–714.
- REED, C. F. 1986. *Flora of the Serpentinite Formations in Eastern North America, with Descriptions of the Geomorphology and Mineralogy at the Formations*. Contrib. 30. Reed Herbarium, Baltimore, MD.
- RICHARDSON, D. H. S. 1995. Metal uptake in lichens. *Symbiosis* 18: 119–127.

- AND R. P. CAMERON. 2004. Cyanolichens: Their response to pollution and possible management strategies for their conservation in northeastern North America. *N. E. Naturalist* 11: 1–22.
- ROBERTS, B. A. AND J. PROCTOR, eds. 1992. *The Ecology of Areas with Serpentinized Rocks: A World View*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- SAFFORD, H. D., J. H. VIERS, AND S. P. HARRISON. 2005. Serpentine endemism in the California flora: A database of serpentine affinity. *Madroño* 52: 222–257.
- SHIMIZU, A. 2004. Community structure of lichens in the volcanic highlands of Mt. Tokachi, Hokkaido, Japan. *Bryologist* 107: 141–151.
- SIROIS, L., F. LUTZONI, AND M. M. GRANDTNER. 1988. Les lichens sur serpentine et amphibolite du plateau du Mont Albert, Gaspésie, Québec. *Canad. J. Bot.* 66: 851–862.
- SULLIVAN, T. J. 1996. *The lichens of Acadia National Park, Maine*. Ph.D. dissertation, University of Minnesota, Minneapolis, MN.
- WERNER, R. G. 1956. Etudes ecologiques sur les lichens des terrains schisteux maritimes. *Bull. Soc. Sci. Nancy*, n. ser. 15: 137–152.
- WILSON, M. J. 1995. Interactions between lichens and rocks: A review. *Cryptog. Bot.* 5: 299–305.

APPENDIX
LICHENS REPORTED FROM PINE HILL AND CALLAHAN MINE

Lichens reported from Pine Hill and Callahan Mine, including taxa listed in Harris et al. (2007) and Rajakaruna et al. (2011). * = new site reports, with substrate indicated by subscripts “r” for rock and “s” for soil. Callahan Mine taxa include a report of where they were collected: TP = tailings pond, WR = waste rock piles, GP = at shore of Goose Pond, IB = in-between. Literature using older nomenclature was interpreted with Esslinger (2012). Mont Albert = reported from Mont Albert (Québec) serpentine in Sirois et al. (1988); Acadia NP = reported from Acadia National Park in Sullivan 1996; X = present, – = absent. Superscripts: ¹metal-tolerant species (Plantlife 2012); ²previous authors may not have distinguished between subspecies or chemospecies. Nomenclature and author citations follow Index Fungorum Partnership (2014).

Species	Pine	Callahan Mine				Mont		Acadia	
	Hill	TP	WR	GP	IB	Albert	NP	NP	
<i>Acarospora fuscata</i> (Nyl.) Th. Fr.	X	–	–	X	–	–	–	X	
<i>Acarospora sinopica</i> (Wahlenb.) Körb. ¹	–	–	X	–	–	–	–	–	
<i>Amanidinea punctata</i> aggr. (Hoffm.) Coppins & Scheid.	*X	–	X	X	–	–	–	X	
<i>Anaptychia palmulata</i> (Michx.) Vain.	X	–	–	–	–	–	–	X	
<i>Aspicilia cinerea</i> (L.) Körb.	–	–	–	X	–	–	–	X	
<i>Aspicilia</i> aff. <i>verrucigera</i> Hue	–	–	X	–	–	–	–	X	
<i>Baeomyces rufus</i> (Huds.) Rebert.	–	–	–	X	–	X	–	X	
<i>Buellia lepidastrata</i> (Tuck.) Tuck.	*X _r	–	–	–	–	–	–	–	
<i>Buellia ocellata</i> (Flörke ex Flot.) Körb.	X	–	X	–	–	–	–	–	
<i>Caloplaca holocarpa</i> (Hoffm.) A.E. Wade	*X _r	–	X	X	–	X	–	X	
<i>Caloplaca lithophila</i> H. Magn.	X	–	–	–	–	–	–	–	
<i>Caloplaca microthallina</i> Wedd.	X	–	–	–	–	–	–	X	
<i>Caloplaca scopularis</i> (Nyl.) Lettau	X	–	–	–	–	–	–	X	
<i>Candelariella aurella</i> (Hoffm.) Zahlbr.	X	–	X	X	–	–	–	X	
<i>Candelariella vitellina</i> (Hoffm.) Müll. Arg.	X	–	X	X	–	X	–	X	
<i>Catillaria lenticularis</i> (Ach.) Th. Fr.	X	–	–	–	–	–	–	–	

APPENDIX. CONTINUED.

Species	Pine Hill	Callahan Mine			Mont Albert	Acadia NP
		TP	WR	GP		
<i>Cladonia acuminata</i> (Ach.) Norri.	X	-	-	-	X	-
<i>Cladonia arbuscula</i> subsp. <i>mitis</i> (Sandst.) Ruoss (syn. <i>Cladonia mitis</i>)	X	-	-	-	X	X
<i>Cladonia boryi</i> Tuck.	X	-	-	-	-	X
<i>Cladonia cariosa</i> (Ach.) Spreng.	X	-	-	-	-	-
<i>Cladonia cervicornis</i> subsp. <i>verticillata</i> (Hoffm.) Ahti	-	-	X	-	-	-
<i>Cladonia chlorophaea</i> (Flörke ex Sommerf.) Spreng. ²	*X _s	-	X	-	X	X
<i>Cladonia cristatella</i> Tuck.	X	-	X	-	-	X
<i>Cladonia cryptochlorophaea</i> Asahina ²	*X _s	-	-	-	-	X
<i>Cladonia dimorphocladata</i> Robbins	*X _s	-	-	-	-	-
<i>Cladonia floerkeana</i> (Fr.) Flörke	-	-	-	X	-	X
<i>Cladonia furcata</i> (Huds.) Schrad.	-	-	-	-	X	X
<i>Cladonia gracilis</i> (L.) Willd. subsp. <i>gracilis</i>	-	-	-	-	-	-
<i>Cladonia gracilis</i> subsp. <i>turbinata</i> (Ach.) Ahti	-	-	X	-	-	X
<i>Cladonia grayi</i> G. Merr. ex Sandst. ²	*X _s	-	-	-	-	X
<i>Cladonia macilentata</i> Hoffm.	X	-	-	-	X	X
<i>Cladonia maxima</i> (Asahina) Ahti	-	-	-	X	X	X
<i>Cladonia phyllophora</i> Erhr. ex Hoffm.	-	-	-	X	X	X
<i>Cladonia pleurota</i> (Flörke) Schaer.	X	-	-	-	X	X
<i>Cladonia pocillum</i> (Ach.) O.J. Rich.	-	-	-	-	-	-
<i>Cladonia pixidata</i> (L.) Hoffm.	X	-	X	-	X	X
<i>Cladonia rangiferina</i> (L.) Weber ex F.H. Wigg.	X	-	-	-	X	X
<i>Cladonia rei</i> Schaer.	X	-	X	-	-	-

APPENDIX. CONTINUED.

Species	Pine	Callahan Mine			Mont	Acadia
	Hill	TP	WR	GP	Albert	NP
<i>Cladonia scabruscula</i> (Delise) Leight	-	-	X	X	-	X
<i>Cladonia squamosa</i> (Scop.) Hoffm.	X	-	-	-	X	X
<i>Cladonia stygia</i> (Fr.) Ruoss	-	-	-	X	-	X
<i>Cladonia subcariosa</i> Nyl. (syn. <i>C. polycarpoides</i>)	X	-	-	X	-	-
<i>Cladonia symphyocarpia</i> (Flörke) Fr.	X	-	-	-	-	-
<i>Cladonia turgida</i> Ehrh. ex Hoffm.	X	-	-	-	X	X
<i>Cladonia uncialis</i> (L.) Weber ex F.H. Wigg.	X	-	-	-	X	X
<i>Coccocarpia palmicola</i> (Spreng.) Arv. & D. J. Galloway	*X _r	-	-	-	-	-
<i>Collema subflaccidum</i> Degel.	X	-	-	-	-	X
<i>Dermatocarpon leptophylloides</i> (Nyl.) Zahlbr.	*X _r	-	-	-	-	-
<i>Dermatocarpon minutum</i> (L.) W. Mann	X	-	-	-	-	-
<i>Dibaeis baomyces</i> (L. f.) Rambold & Hertel	X	-	-	X	-	X
<i>Diploschistes muscorum</i> (Scop.) R. Sant.	-	-	X	X	-	-
<i>Evernia mesomorpha</i> Nyl.	-	-	-	X	-	X
<i>Flavoparmelia caperata</i> (L.) Hale	X	-	-	-	-	X
<i>Fuscidea pusilla</i> Tonsberg	-	-	-	X	-	-
<i>Fuscopannaria praetermissa</i> (Nyl.) P.M. Jørg.	X	-	-	-	-	-
<i>Hypogymnia physodes</i> (L.) Nyl.	-	-	X	X	X	X
<i>Lecanora argentea</i> Oxner & Volkova	X	-	-	-	-	-
<i>Lecanora dispersa</i> (Pers.) Rohl.	X	-	X	X	-	-
<i>Lecanora polytropa</i> (Ehrh.) Rabenh.	X	-	X	X	X	X
<i>Lecidea auriculata</i> Th. Fr.	-	-	*X _r	-	-	-
<i>Lecidella patavina</i> (A. Massal.) Knoph & Leuckert	*X _r	-	-	-	-	-
<i>Lecidella stigmatea</i> (Ach.) Hertel & Leuckert	X	-	-	-	X	-

APPENDIX. CONTINUED.

Species	Pine		Callahan Mine				Mont Albert	Acadia NP
	Hill	TP	WR	GP	IB			
<i>Peltigera rufescens</i> (Weiss) Humb.	X	-	X	X	-	-	-	
<i>Pertusaria amara</i> (Ach.) Nyl.	X	-	-	-	-	-	X	
<i>Phaeophyscia adiantola</i> (Essl.) Essl.	X	-	-	-	-	-	X	
<i>Phaeophyscia rubropulchra</i> (Degel.) Moberg	X	-	-	-	-	-	X	
<i>Phaeophyscia sciastra</i> (Ach.) Moberg	X	-	-	X	-	-	X	
<i>Physcia adscendens</i> (Fr.) H. Olivier	-	-	-	X	-	-	X	
<i>Physcia aipolia</i> (Ehrh. ex Humb.) Hampe ex Fürnr.	-	-	-	X	-	X	X	
<i>Physcia caesia</i> (Hoffm.) Fürnr.	X	-	-	-	-	X	X	
<i>Physcia dubia</i> (Hoffm.) Lettau	-	-	X	X	-	-	X	
<i>Physcia phaea</i> (Tuck.) J.W. Thomson	-	-	-	X	-	-	X	
<i>Physcia subtilis</i> Degel.	-	-	-	X	-	-	X	
<i>Physcia tenella</i> (Scop.) DC.	*X _r	-	-	-	-	-	X	
<i>Placidium squamulosum</i> (Ach.) Breuss	*X _r	-	-	-	-	-	X	
<i>Placynthiella hyporhoda</i> (Th. Fr.) Coppins & P. James ¹	-	-	-	*X _s	-	-	-	
<i>Placynthiella icmalea</i> (Ach.) Coppins & P. James	X	-	-	X	-	-	X	
<i>Placynthiella uliginosa</i> (Schrad.) Coppins & P. James	*X _s	-	-	-	-	X	-	
<i>Porocyphus coccodes</i> Flot. ex Körb.	*X	-	-	-	-	-	-	
<i>Porpidia crustulata</i> (Ach.) Hertel & Knoph	-	-	X	-	-	X	X	
<i>Porpidia macrocarpa</i> (DC.) Hertel & A.J. Schwab	-	-	-	X	-	X	X	
<i>Porpidia subsimplex</i> (H. Magn.) Fryday	X	-	-	-	-	-	X	
<i>Psorula rufonigra</i> (Tuck.) Gotth. Schneid.	X	-	-	-	-	-	-	
<i>Punctelia rufecta</i> (Ach.) Krog	-	-	-	X	-	-	X	
<i>Pyrenocarpon thelostomum</i> (Ach. ex J. Harriman) Coppins & Aptroot	*X _r	-	-	-	-	-	-	

APPENDIX. CONTINUED.

Species	Pine	Callahan Mine			Mont Albert	Acadia NP
	Hill	TP	WR	GP		
<i>Ramalina intermedia</i> (Delise ex Nyl.) Nyl.	-	-	-	X	-	X
<i>Rhizocarpon cinereovirens</i> (Müll. Arg.) Vain.	-	-	-	X	X	X
<i>Rhizocarpon dispersum</i> (Nägeli ex Hepp) Müll. Arg.	*X _r	-	-	-	-	-
<i>Rhizocarpon distinctum</i> Th. Fr.	-	-	*X _r	-	-	-
<i>Rhizocarpon geminatum</i> Körb.	X	-	-	-	-	-
<i>Rhizocarpon grande</i> (Flörke ex Flot.) Arnold	-	-	X	X	-	X
<i>Rhizocarpon infernulum</i> (Nyl.) Lynge	-	-	-	X	-	-
<i>Rhizocarpon lecanorinum</i> Anders	-	-	X	X	-	X
<i>Rhizocarpon reductum</i> Th. Fr. (syn. <i>Rhizocarpon obscuratum</i>)	X	-	X	X	-	X
<i>Rhizocarpon rubescens</i> Th. Fr.	-	-	X	X	-	X
<i>Rinodina gennarii</i> Bagl.	-	-	X	X	-	-
<i>Sarcosagium campestre</i> (Fr.) Poetsch & Schied. ¹	-	*X _s	-	-	-	-
<i>Scoliosporium umbrinum</i> (Ach.) Arnold	X	-	X	X	X	X
<i>Spilonema revertens</i> Nyl.	X	-	-	-	-	-
<i>Steinia geophana</i> (Nyl.) Stein ¹	-	*X _s	-	-	-	-
<i>Stereocaulon condensatum</i> Hoffm. ¹	-	-	-	-	*X _s	-
<i>Stereocaulon dactylophyllum</i> Flörke ¹	-	X	X	X	X	X
<i>Stereocaulon glaucescens</i> Tuck.	X	-	-	-	X	-
<i>Stereocaulon pileatum</i> Ach. ¹	-	-	X	X	-	X
<i>Stereocaulon saxatile</i> H. Magn.	-	-	-	X	-	X
<i>Stereocaulon subcoralloides</i> Nyl.	-	-	*X _s	-	-	X
<i>Stereocaulon tomentosum</i> Th. Fr.	-	-	-	X	X	X
<i>Stigmidium</i> sp. [NBM: TH 283-2]	-	-	-	X	-	-

APPENDIX. CONTINUED.

Species	Pine Hill	Callahan Mine			Mont Albert	Acadia NP
		TP	WR	GP		
<i>Trapelia placodioides</i> Coppins & P. James	-	-	X	-	-	X
<i>Trapeliopsis granulosa</i> (Hoffm.) Lumbsch	*X	X	X	-	X	X
<i>Verrucaria muralis</i> Ach.	-	X	X	-	-	-
<i>Vezeada acicularis</i> Coppins ¹	-	-	*X _s	-	-	-
<i>Xanthoparmelia conspersa</i> (Ehrh. ex Ach.) Hale	*X _r	-	X	-	-	X
<i>Xanthoparmelia cumberlandia</i> (Gyeln.) Hale	X	-	X	-	-	X
<i>Xanthoparmelia plititii</i> (Gyeln.) Hale	X	-	X	-	-	X
<i>Xanthoparmelia viridolubrina</i> (Gyeln.) Lendemer	*X _r	-	X	-	-	-
<i>Xanthoria elegans</i> (Link) Th. Fr.	X	-	X	-	X	X
<i>Xanthoria parietina</i> (L.) Beltr.	X	-	X	-	-	X