1-1-1987

Fluctuating Hydrodynamics in a Dilute Gas

Alejandro Garcia

San Jose State University, alejandro.garcia@sjsu.edu

G. Lie

IBM Corporation

E. Clementi

IBM Corporation

Follow this and additional works at: https://scholarworks.sjsu.edu/physics_astron_pub

Part of the [Other Astrophysics and Astronomy Commons](#), and the [Other Physics Commons](#)

Recommended Citation

Alejandro Garcia, G. Lie, and E. Clementi. "Fluctuating Hydrodynamics in a Dilute Gas" *Physical Review Letters* (1987): 874-877. <https://doi.org/http://dx.doi.org/10.1103/PhysRevLett.58.874>

This Article is brought to you for free and open access by the Physics and Astronomy at SJSU ScholarWorks. It has been accepted for inclusion in Faculty Publications by an authorized administrator of SJSU ScholarWorks. For more information, please contact scholarworks@sjsu.edu.

Fluctuating Hydrodynamics in a Dilute Gas

M. Malek Mansour,^(a) Alejandro L. Garcia,^(a) George C. Lie, and Enrico Clementi

IBM Corporation, Kingston, New York 12401

(Received 26 June 1986)

Hydrodynamic fluctuations in a dilute gas subjected to a constant heat flux are studied by both a computer simulation and the Landau-Lifshitz formalism. The latter explicitly incorporates the boundary conditions of the finite system, thus permitting quantitative comparison with the former. Good agreement is demonstrated.

PACS numbers: 47.90.+a, 05.40.+j, 05.70.Ln

Fluctuating hydrodynamics is a stochastic formulation of standard fluid mechanics.¹ Spontaneous fluctuations of hydrodynamic variables are introduced into the transport equations by the addition of random components to the pressure and heat fluxes. Since these fluxes are not conserved quantities, the correlations between the random components are expected to be short ranged and short lived so that at hydrodynamic scales they are assumed to be Dirac-delta correlated. Their strengths are then chosen to yield the correct equilibrium thermodynamic fluctuations as derived from the Gibbs distribution. Nowadays there are various ways to derive the Landau-Lifshitz fluctuating hydrodynamics and there is general agreement about its validity, at least in near-equilibrium situations.²

Extension of the theory to nonequilibrium systems leads to predictions of the asymmetry of the Brillouin lines in a liquid subjected to a constant heat flux.³⁻⁶ Kinetic theory provides further support for these predictions.^{7,8} Although these theoretical results are in agreement with light-scattering experiments,^{9,10} the importance of the nonlinearities¹¹ and the influence of the boundaries¹² remain under discussion (see also the work of Tremblay¹³). In any case, the question arises as to the applicability of the fluctuating-hydrodynamics formalism to systems under strong nonequilibrium constraints. One way to address these questions is through particle simulations.

In this article we study a dilute hard-sphere gas bounded by two parallel plates located at $y=0$ and $y=L$,

using both the fluctuating-hydrodynamics formalism and a Boltzmann Monte Carlo particle simulation. The plates act as infinite reservoirs so that by fixing their temperatures one can impose the desired temperature gradient across the system. As can be checked easily from the macroscopic hydrodynamic equations, the heat flux in the stationary state is constant and the velocity is zero [note that there is no instability because we do not include external fields (gravity) in our formulation¹⁴]. To study the fluctuations, we first linearize the fluctuating-hydrodynamics equations around the macroscopic stationary state. Since we are mainly interested in the influence of nonequilibrium constraints and since the particle simulations with which we compare our results employ periodic boundary conditions in the x and z directions, we shall limit ourselves to reduced quantities, defined as

$$\delta A(y) \equiv \frac{1}{S} \int_0^{L_x} dx \int_0^{L_z} dz \delta A(x, y, z), \quad (1)$$

where A is any dynamical variable and $S \equiv L_x L_z$ is the wall cross section (note that the reduced variables are in fact the zero-wave-vector values of the "parallel" Fourier components of the dynamical variables). It is easy to check that the reduced equations for the x and z components of the velocity fluctuations decouple from the rest and are not influenced by the constraint. We will therefore concentrate our attention on the remaining equations for the reduced mass density $\delta\rho$, the y component of velocity δv , and the temperature δT , which turn out to be

$$\partial \delta \rho / \partial t = -\partial \rho_0 \delta v / \partial y, \quad (2)$$

$$\rho_0 \frac{\partial \delta v}{\partial t} = -R \frac{\partial}{\partial y} (T_0 \delta \rho + \rho_0 \delta T) + \frac{4}{3} \frac{\partial \eta_0}{\partial y} \frac{\partial \delta v}{\partial y} - \frac{\partial s_{yy}}{\partial y}, \quad (3)$$

$$\frac{3}{2} \rho_0 R \frac{\partial \delta T}{\partial t} = -\frac{3}{2} R \rho_0 \delta v \frac{\partial T_0}{\partial y} - P_0 \frac{\partial \delta v}{\partial y} + \frac{\partial}{\partial y} \left(\delta \kappa \frac{\partial T_0}{\partial y} + \kappa_0 \frac{\partial \delta T}{\partial y} \right) - \frac{\partial g_y}{\partial y}, \quad (4)$$

where the subscript 0 indicates *local* macroscopic quantities, R the Boltzmann constant divided by the mass, η_0 the shear viscosity, κ_0 the thermal conductivity, and P_0 the pressure. s_{yy} and g_y are the random components of the pressure and heat fluxes, respectively, with the following covariances¹:

$$\langle s_{y,y}(y, t) s_{y,y}(y', t') \rangle = \frac{8}{3} k_B T_0 (\eta_0 / S) \delta(y - y') \delta(t - t'), \quad (5a)$$

$$\langle g_y(y, t) g_y(y', t') \rangle = 2k_B T_0^2 (\kappa_0 / S) \delta(y - y') \delta(t - t'), \quad \langle s_{y,y}(y, t) g_y(y', t') \rangle = 0. \quad (5b,c)$$

In writing Eqs. (2)–(4) we have made use of the closure relations for a dilute gas, $P(\rho, T) = R\rho T$ and $e(\rho, T) = 3\rho RT/2$ where e is the internal energy density. If the force between the particles is purely repulsive and obeys a power law, then the transport coefficients are only functions of temperature as¹⁵ $\eta_0 = \eta_c T_0^\alpha$ and $\kappa_0 = \kappa_c T_0^\alpha$, so that in Eq. (4) we can set $\delta\kappa = \alpha\kappa_0\delta T/T_0$. For a hard-sphere gas, the exponent α is $\frac{1}{2}$ and, from Chapman-Enskog theory, $\kappa_c/\eta_c = 15R/4$.

There remains the problem of specifying the boundary conditions for Eqs. (2)–(4). If we assume that the state of the walls is statistically independent with respect to the system, then the boundary conditions for δT are

$$\delta T(y=0, t) = \delta T(y=L, t) = 0. \quad (6)$$

The boundary conditions for δv follow from the conservation of the total particle number; the continuity equation yields

$$\rho_0(y)\delta v(y)|_{\text{boundaries}} = 0. \quad (7)$$

This completes our specification of the boundary conditions.

It may seem strange that we do not have to specify any boundary conditions for $\delta\rho$. From a physical point of view, this comes from the fact that the state of the wall can only constrain the temperature and velocity of the gas at the wall, whereas the behavior of the density close to the wall is entirely determined by the internal dynamics of the system. From the mathematical point of view, it can be shown that for any given initial condition $\delta\rho(y, 0)$, $\delta\mathbf{u}(y, 0)$, $\delta T(y, 0)$, the boundary conditions for $\delta\mathbf{u}$ and δT are sufficient to specify completely the solution of the system.¹⁶

Because the coefficients and the noise are both space dependent and because we are dealing with a finite system, it is no longer possible to use elementary transform methods to solve the above fluctuating-hydrodynamics equations (for large enough systems, an expansion in the wave number of the gradient can still be used^{4,17}). An alternative approach would be a direct computer simulation of the Langevin equations (2)–(4). This is a useful method in the study of fluctuating-hydrodynamics equations in two or three dimensions.¹⁸ Here, instead, we construct the evolution equations for the correlation functions and numerically solve these equations by relaxation.¹⁹ The first step in this approach is the evaluation of the static correlation functions, which can then be used to compute the dynamical correlation functions as an initial value problem. In this short Letter we deal only the static correlation functions.

To proceed further, we appeal to a very useful identity of the theory of stochastic processes²⁰, namely, given that

$$dc_i/dt = f_i(c_1, \dots, c_n) + F_i(t), \quad i = 1, 2, \dots, n, \quad (8a)$$

where the f_i 's are arbitrary analytic functions of c_i 's, and

$F_i(t)$'s are multi-Gaussian white-noise processes with covariances

$$\langle F_i(t)F_j(t') \rangle = Q_{ij}\delta(t-t'), \quad (8b)$$

then

$$\langle c_i(t)F_j(t') \rangle \equiv \begin{cases} \frac{1}{2} Q_{ij}, & t = t', \\ 0, & t < t'. \end{cases} \quad (8c)$$

For finite n , this identity is easily proved by writing the Fokker-Planck equation corresponding to (8a) and from it deriving the second-moment equations. A comparison with the second-moment equations derived directly from (8a) then leads to the relation (8c). These relations remain valid for $n \rightarrow \infty$ although, from a strictly mathematical point of view, some special care is needed in the continuum case. Using the relation (8c), one can derive the evolution equations for the equal-time correlation functions. If one discretizes the spatial derivatives, the steady-state solution may then be obtained by relaxation methods.

The primary purpose of this work is to compare the predictions of the fluctuating-hydrodynamics theory with particle-simulation results (the nonequilibrium effects we are considering are too subtle to be readily studied by laboratory experiments). Molecular-dynamics simulations prove to be too slow and have, thus far, yielded only qualitative results.²¹ We rely, instead, on results obtained by a Boltzmann Monte Carlo simulation originally developed for rarefied gas studies by Bird²² (some limitations of this method are discussed by Meiburg²³). We have considered a system containing 20000 particles between two thermal plates 50 mean free paths (λ) apart and held at different temperatures. Here we report the results for a temperature gradient of 0.04° per mean free path. Distances and velocities are scaled by λ and the most probable speed, $(2k_B T/m)^{1/2}$, respectively; the

FIG. 1. Comparison of the temperature-temperature spatial correlation functions obtained from the fluctuating-hydrodynamics formalism (solid curve) and Boltzmann Monte Carlo particle simulation. Note that the local equilibrium delta-function contribution has been removed.

FIG. 2. Comparison of the density-velocity spatial correlation functions obtained from the fluctuating-hydrodynamics formalism (solid curve) and Boltzmann Monte Carlo particle simulation.

mass is set equal to 1 and k_B to $\frac{1}{2}$ (see the work of Garcia²⁴ for details). The statistical error is estimated to be about 10% for the temperature autocorrelation function (Fig. 1) and less than 5% for the other correlation functions (Figs. 2 and 3). The local equilibrium contributions to the correlation functions are removed and, as a result, larger errors are to be expected at the central peak. We note that there are no free parameters in the analysis; the solution of the correlation-function equations is entirely specified once the simulation parameters are given. These are, in the reduced units, $T(y=0)=1$, $T(y=L)=3$, and $\eta_c=5\pi^{1/2}\rho_{eq}/16$, where ρ_{eq} is N/L . Because of a small slip in the temperature profile, $T(y=L)$ is set equal to 2.95 in the hydrodynamic equations (2)–(4).

The program was run in parallel on two FPS264 array processors attached to the 1CAP2 system at IBM Kingston for 2×10^5 collisions per article. Figure 1 shows the temperature-temperature static correlation function which is clearly long ranged. Despite some statistical scatter, the simulation results show quite good agreement with the fluctuating-hydrodynamics results. The nonequilibrium contribution to the global temperature fluctuation (defined as the space average of the static temperature autocorrelation function) is found to be proportional to the square of the temperature gradient. Further studies with different system sizes indicate that for fixed temperature gradient it increases with the length of the system. These observations are in agreement with previous work on model systems.²⁵ In Figs. 2 and 3, we depict the density-velocity and density-density static correlation functions, respectively. Because of the conservation of the total mass, the static density autocorrelation function is strictly negative; its space integral compensates exactly for the local equilibrium contribution. Both curves show much better agreement with the fluctuating-hydrodynamics predictions. Similar agree-

FIG. 3. Comparison of the density-density spatial correlation functions obtained from the fluctuating-hydrodynamics formalism (solid curve) and Boltzmann Monte Carlo particle simulation. Note that the local equilibrium delta-function contribution has been removed.

ment is found for all the static correlation functions investigated. For completeness, we are also studying the dynamic correlation functions and are experimenting with molecular-dynamics simulations for dense systems using more realistic interaction potentials.

Our present observations suggest that the fluctuating-hydrodynamics equations are valid at length scales of a few mean free paths even in the presence of strong non-equilibrium constraints, at least for a dilute gas (see also Alder and Wainwright²⁶). Had the data shown otherwise then a strictly microscopic formulation in kinetic theory would have been the only recourse. We consider this a fortunate development which will encourage future work in this direction. For instance, recent large-scale molecular-dynamics results demonstrate the feasibility of the observation of macroscopic hydrodynamic phenomena such as vortex formation and shedding past an obstacle^{23,27,28} in particle simulations. The next major step, of course, will be the study of fluctuations near hydrodynamic instabilities by computer simulations. Our results indicate that fluctuating hydrodynamics provides a promising way to tackle this problem, at least before and, probably, close to the instability.¹⁴

The authors wish to thank Prof. G. Nicolis, Prof. I. Prigogine, Dr. L. Hannon, Dr. F. Baras, Dr. J. W. Turner, and Dr. M. Mareschal for their stimulating discussions and helpful suggestions. Two of us (A.G. and M.M.) also wish to thank the Computational Support staff, particularly Dr. R. Caltabiano and Dr. M. Russo, of IBM Kingston for their assistance of 1CAP2.

(a)Permanent address: Service de Chimie Physique II,

Université Libre de Bruxelles, Campus Plaine, C.P. 231, 1050 Bruxelles, Belgium.

¹L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon Press, Oxford, 1959).

²For a review see, for instance, M. H. Ernst and E. G. D. Cohen, *J. Stat. Phys.* **25**, 153 (1991).

³I. Procaccia, D. Ronis, and I. Oppenheim, *Phys. Rev. Lett.* **42**, 287 (1979); D. Ronis, I. Procaccia, and I. Oppenheim, *Phys. Rev. A* **19**, 1324 (1979); D. Ronis and S. Putterman, *Phys. Rev. A* **22**, 773 (1980).

⁴A.-M. S. Tremblay, M. Arai, and E. Siggia, *Phys. Rev. A* **23**, 1451 (1981).

⁵G. Van Der Zwan, D. Bedaux, and P. Mazur, *Physica (Amsterdam)* **107A**, 491 (1981).

⁶T. R. Kirkpatrick, E. G. D. Cohen, and J. R. Dorfman, *Phys. Rev. Lett.* **44**, 472 (1980).

⁷T. R. Kirkpatrick, E. G. D. Cohen, and J. R. Dorfman, *Phys. Rev. Lett.* **42**, 862 (1979).

⁸J. Dufty, in *Spectral Line Shapes*, edited by P. Wende (de Gruyter, Berlin, 1981), p. 1143.

⁹D. Beysens, Y. Garrabos, and G. Zaleger, *Phys. Rev. Lett.* **45**, 403 (1980); D. Beysens, *Physica (Amsterdam)* **118A**, 250 (1983).

¹⁰R. Penney, H. Kiefte, and J. M. Clouter, *Bull. Can. Assoc. Phys.* **39**, BB8 (1983).

¹¹T. R. Kirkpatrick and E. G. D. Cohen, *Phys. Lett.* **78A**, 350 (1980); D. Ronis and I. Procaccia, *Phys. Rev. A* **26**, 1812 (1982).

¹²G. Satten and D. Ronis, *Phys. Rev. A* **26**, 940 (1982); T. R. Kirkpatrick and E. G. D. Cohen, *Phys. Rev. A* **26**, 972 (1982).

¹³A nice review of the field is given by A. M. Tremblay, in *Recent Developments in Nonequilibrium Thermodynamics*, edited by J. Casas-Vasquez, D. Jou, and G. Lebon (Springer-

Verlag, Berlin, 1984).

¹⁴R. Schmitz and E. G. D. Cohen, *J. Statist. Phys.* **38**, 285 (1985), and **40**, 431 (1985).

¹⁵P. Resibois and M. DeLeener, *Classical Kinetic Theory of Fluids* (Wiley and Sons, New York, 1977).

¹⁶M. Malek Mansour, J. W. Turner, A. L. Garcia, and G. Nicolis, to be published.

¹⁷I. Procaccia, D. Ronis, and I. Oppenheim, *Phys. Rev. A* **20**, 2533 (1979); D. Ronis, I. Procaccia, and J. Machata, *Phys. Rev. A* **22**, 714 (1980); A. Perez-Madrid and J. M. Rubi, *Phys. Rev. A* **33**, 2716 (1986).

¹⁸A. L. Garcia, M. Malek Mansour, G. Lie, and E. Clementi, to be published.

¹⁹D. Potter, *Computational Physics* (John Wiley and Sons, New York, 1979).

²⁰L. Arnold, *Stochastic Differential Equations: Theory and Applications* (Wiley, New York, 1974); J. W. Dufty, J. J. Brey, and M. C. Marchetti, *Phys. Rev. A* **33**, 4307 (1986).

²¹M. Mareschal and E. Kestemont, *Phys. Rev. A* **30**, 1158 (1984).

²²G. Bird, *Molecular Gas Dynamics* (Clarendon, Oxford, 1976).

²³W. Meiburg, Deutschen Forschungs- und Versuchsanstalt für Luft- und Raumfahrt (DFVLR) Report No. FB85-13, 1985 (unpublished).

²⁴A. Garcia, *Phys. Rev. A* **34**, 1454 (1986).

²⁵G. Nicolis and M. Malek Mansour, *Phys. Rev. A* **29**, 2845 (1984).

²⁶B. J. Alder and T. E. Wainwright, *Phys. Rev. A* **1**, 18 (1970).

²⁷L. Hannon and E. Clementi, to be published.

²⁸D. Rappaport and E. Clementi, *Phys. Rev. Lett.* **57**, 695 (1986).