

Fall 2012

Dynamics of *Vibrio* with virulence genes detected in Pacific harbor seals (*Phoca vitulina richardii*) off California: implications for marine mammal health

Stephanie Nichole Hughes
San Jose State University

Follow this and additional works at: https://scholarworks.sjsu.edu/etd_theses

Recommended Citation

Hughes, Stephanie Nichole, "Dynamics of *Vibrio* with virulence genes detected in Pacific harbor seals (*Phoca vitulina richardii*) off California: implications for marine mammal health" (2012). *Master's Theses*. 4235.

DOI: <https://doi.org/10.31979/etd.5wc5-enkg>

https://scholarworks.sjsu.edu/etd_theses/4235

This Thesis is brought to you for free and open access by the Master's Theses and Graduate Research at SJSU ScholarWorks. It has been accepted for inclusion in Master's Theses by an authorized administrator of SJSU ScholarWorks. For more information, please contact scholarworks@sjsu.edu.

DYNAMICS OF *VIBRIO* WITH VIRULENCE GENES DETECTED IN PACIFIC
HARBOR SEALS (*PHOCA VITULINA RICHARDII*) OFF CALIFORNIA:
IMPLICATIONS FOR MARINE MAMMAL HEALTH

A Thesis

Presented to

The Faculty of Moss Landing Marine Laboratories

San José State University

In Partial Fulfillment

of the Requirements for the Degree

Master of Science

by

Stephanie N. Hughes

December 2012

© 2012

Stephanie N. Hughes

ALL RIGHTS RESERVED

The Designated Thesis Committee Approves the Thesis Titled

DYNAMICS OF *VIBRIO* WITH VIRULENCE GENES DETECTED IN PACIFIC
HARBOR SEALS (*PHOCA VITULINA RICHARDII*) OFF CALIFORNIA:
IMPLICATIONS FOR MARINE MAMMAL HEALTH

by

Stephanie N. Hughes

APPROVED FOR MOSS LANDING MARINE LABORATORIES

SAN JOSÉ STATE UNIVERSITY

December 2012

Dr. James T. Harvey

Moss Landing Marine Laboratories

Dr. Michael Graham

Moss Landing Marine Laboratories

Dr. Frances M.D. Gulland

The Marine Mammal Center

ABSTRACT

DYNAMICS OF *VIBRIO* WITH VIRULENCE GENES DETECTED IN PACIFIC HARBOR SEALS (*PHOCA VITULINA RICHARDII*) OFF CALIFORNIA: IMPLICATIONS FOR MARINE MAMMAL HEALTH

by Stephanie N. Hughes

Given their coastal site fidelity and opportunistic foraging behavior, harbor seals (*Phoca vitulina*) may serve as sentinels for coastal ecosystem health. Seals using urbanized coastal habitat can acquire enteric bacteria, including *Vibrio*, that may affect their health. To understand *Vibrio* dynamics in seals, demographic and environmental factors were tested for predicting potentially virulent *Vibrio* in free-ranging and stranded Pacific harbor seals (*P. v. richardii*) off the coast of California. *Vibrio* prevalence did not vary with season and was greater in free-ranging seals (29%, $n = 319$) compared with stranded seals (17%, $n = 189$). Of the factors tested, location, turbidity, and/or salinity best predicted *Vibrio* prevalence in free-ranging seals. The relationship of environmental factors with *Vibrio* prevalence differed by location and may be related to oceanographic or terrestrial contributions to water quality. *Vibrio parahaemolyticus*, *V. alginolyticus*, and *V. cholerae* were observed in seals with *V. cholerae* found almost exclusively in stranded pups and yearlings. Additionally, virulence genes (*trh* and *tdh*) were detected in *V. parahaemolyticus* isolates. *Vibrio cholerae* isolates lacked targeted virulence genes, but were hemolytic. Three out of four stranded pups with *V. parahaemolyticus* (*trh*+, and/or *tdh*+) died in rehabilitation, but the role of *Vibrio* in causing mortality is unclear, and *Vibrio* expression of virulence genes should be investigated. Considering that

humans share the environment and food resources with seals, potentially virulent *Vibrio* observed in seals also may be of concern to human health.

ACKNOWLEDGEMENTS

I consider my experience at Moss Landing Marine Labs a memorable adventure, and this thesis a cherished accomplishment. This work was made possible by the guidance, support, and comedic relief of many individuals. With immense gratitude, I would like to thank my advisor and friend, Dr. Jim Harvey. Jim, you provided for me the opportunity to pursue the work I love, challenged my academic abilities, sharpened my analytical skills, and shared valuable perspective on life outside MLML. Your wit and humor kept me entertained and calm in the lab and field. It was greatly appreciated. Admittedly, I was amazed and envious that you had energy to spare after we spent hours working up dozens of seals on a “good” Tomales day. I also would like to thank my committee members, Drs. Frances Gulland and Mike Graham. Frances, you have been an incredible mentor. I am deeply appreciative of your support and encouragement. Thank you for fostering my excitement for this project, entertaining my many tangents, and for being a humble, fun, and brilliant role model. Mike, you lit the spark within me that has forever changed the way I view nature and science. Your enthusiasm for the natural world is radiant, and infectious. Naturally, I am susceptible! Thank you, because that energy stuck with me and was essential for completing this work.

Special thanks to Liz McHuron, Denise Greig, and Jenny Carlson for their unconditional support, encouragement, and constructive criticism throughout this work. I also would like to thank Suzanne Manugian, Scott Hansen, Alex Olson, Deasy Lontoh, and Casey Clark for their efforts and support in the field. Thanks to Tenaya Norris and Tanya Novak for providing valuable Matlab expertise, and Michelle Marraffini for help with Geneious. Thanks to Drs. Barbara Byrne, and Dr. Woutrina Miller, I had a second

home at UCD that provided the workspace, equipment, and expertise for the molecular aspect of this project. Dr. Mark Strom generously donated reference strains that I used as controls. I am also grateful for the efforts and support of countless volunteers, faculty, and staff that provided the resources and manpower to get the job done.

Lastly, I would like to thank my family. My mother, Heidi, instilled in me a sense of adventure. Mom, your zest for life and love of nature is my inspiration. My father, Bob, taught me perseverance. Pops, I can't thank you enough for your love and support throughout my academic career. I am forever indebted to my husband, Brian, who introduced to me the wonder of the marine world and has provided unconditional support, laughter, and love ever since. Of course, I have Uncle Wooly and Grams to thank for their relentless humor, and optimism. Grams, thank you for catching pill bugs with me in your backyard. That was my beginning as a naturalist. Not to mention drinking lemon drops with you at the cabin certainly can solve any *Vibrio* conundrum!

Financial support for sample processing was provided by the Marine Mammal Center and funding for molecular reagents was provided by the Earl and Ethyl Myers Oceanographic and Marine Biology Trust, CSU-COAST Marine Science Research Award, and the Packard Foundation. I also would like to thank the National Parks Service, Don Edwards National Wildlife Refuge, and the Humboldt Bay NWR Complex for supporting this work. Samples were collected under permits issued by NMFS (Nos. 555-1870, 373-1868), USFWS (Nos. 2009-041 2011-002, 10007), NPS (81640-2011-002), and IACUC protocol No. 948 issued by San Jose State University.

Table of Contents

List of Tables.....	viii
List of Figures.....	ix
Introduction.....	1
Materials and Methods.....	3
<i>Isolation and characterization of Vibrio spp. from free-ranging and stranded seals</i>	3
<i>Environmental data collection</i>	8
<i>Data analysis</i>	8
Results.....	10
<i>Vibrio prevalence in free-ranging harbor seals</i>	10
<i>Vibrio prevalence in stranded seals</i>	14
<i>Comparison of Vibrio prevalence, species distribution, and virulence profiles between free-ranging and stranded seals</i>	15
Discussion.....	19
References.....	29
Appendix A. Descriptive statistics, and bin criteria used for categorizing environmental predictor variables to be used in multivariate logistic regression analysis for SFB, from 2007 to 2011.....	38
Appendix B. Descriptive statistics, and bin criteria used for categorizing environmental predictor variables to be used in multivariate logistic regression analysis for ES in 2010.....	38
Appendix C. Sequence confirmation for target genes for a subset of isolates collected from harbor seals.....	39

List of Tables

Table 1. Results from logistic regression testing location as a predictor for *Vibrio* presence and absence in seals sampled in 2007 to 2011.....12

Table 2. Results from multivariate backwards stepwise logistic regression testing environmental variables as predictors of the presence or absence of *Vibrio* in free-ranging seals sampled from Elkhorn Slough in 2010 and then San Francisco Bay from 2007 to 2011.....13

Table 3. Prevalence of *Vibrio* in free-ranging and stranded pups and yearlings positive for *Vibrio* per location from 2007 to 2011.....16

Table 4. Virulence gene profiles for *V. parahaemolyticus* isolates collected from free-ranging seals from Elkhorn Slough (ES), San Francisco Bay (SFB), Tomales Bay (TB), and Humboldt Bay (HB), and stranded seals (TMMC) in 2010 and 2011. Sample sizes represent the number of isolates tested.....19

List of Figures

- Figure 1. Sampling sites and sample sizes for free-ranging harbor seals ($n = 319$, black dots) and area between San Luis Obispo and Mendocino (black bars) where stranded animals were sampled ($n = 189$) during admission to The Marine Mammal Center in 2007 through 2011.....5
- Figure 2. Proportion of free-ranging harbor seals with *Vibrio* (seals confirmed with *Vibrio*/ total seals sampled) by year and location (SFB = San Francisco Bay, TB = Tomales Bay, ES = Elkhorn Slough). Seals were sampled from ES only in 2010. Sample sizes are indicated in parentheses.....11
- Figure 3. Proportion of stranded pups and yearlings admitted to The Marine Mammal Center with *Vibrio*, 2007 to 2011. Sample sizes are indicated in parentheses.....14
- Figure 4. Proportions of *Vibrio* species isolated from (A) free-ranging harbor seals sampled in Humboldt Bay (HB, $n = 13$), Tomales Bay (TB, $n = 20$), San Francisco Bay (SFB, $n = 32$) and Elkhorn Slough (ES, $n = 28$), and (B) harbor seals stranded in TB ($n = 2$), SFB ($n = 11$), and Monterey Bay (MB, $n = 6$). Mixed *Vibrio* Culture are isolates that could not be identified to species level, and samples with more than one species of *Vibrio* detected....17

Introduction

Marine mammals are sensitive to changing environmental conditions and can serve as sentinels for ecosystem health [11, 13, 35, 60]. Long-term monitoring of marine mammal health includes measuring contaminants, trace elements, biotoxins from harmful algal blooms, baseline blood values, and prevalence of infectious disease so that deviations from normal values may be detected [10, 12, 32, 33, 35-38]. Cumulative effects of one or more of these factors may have contributed to the deterioration of marine mammal health [10, 11, 13, 35]. Because infectious marine diseases appear to be increasing and are of immediate concern to marine mammal health, the dynamics and virulence potential of aquatic microbes should be investigated [4, 10, 28, 38, 48, 75].

Baseline epidemiological data on marine pathogens can be obtained from a representative marine mammal to aid in disease mitigation in marine mammals [10, 36, 48, 65, 83]. Harbor seals (*Phoca vitulina*) are an excellent indicator species because they are long-lived and upper-level trophic consumers that inhabit coastal areas throughout the northern hemisphere. Harbor seals often have strong site fidelity to areas near dense human populations (e.g., San Francisco Bay), and forage opportunistically on available benthic and pelagic prey [9, 23, 62, 69, 84, 91]. Terrestrial inputs from urbanized coastal communities may alter the quality of habitat or food resources exploited by harbor seals [32, 53, 58, 70, 72, 82]. These inputs also may contribute to increases in marine pathogens [39, 42], thereby leading to an increasing incidence of disease in marine animals including harbor seals.

Current knowledge of the diversity and ecology of marine pathogens in harbor seals is limited to clinical cases, serological surveys, zoonotic cases (marine mammal caretakers), and epidemics associated with animal-stranding events [10, 37, 38, 73, 78, 83, 92]. The most common cause of live harbor seal strandings in central California during the last ten years was malnutrition (52%) [16]. Malnourished individuals may be more susceptible to enteric pathogens [44]. To better understand impacts of enteric pathogens on the health of harbor seals, enumerating these pathogens in healthy and stranded seals is warranted [10, 16, 32, 48, 51].

Marine pathogens of the genus *Vibrio* are of concern to marine mammal and ecosystem health. *Vibrios* are facultative anaerobes that can be found in aquatic environments throughout the world [18, 24]. Pathogenic strains may proliferate following environmental perturbation, and nonpathogenic strains may become competent pathogens via inter-microbial gene transfer [4, 7, 18, 24, 34, 39, 57, 63, 79]. The diversity and versatility of this group of bacteria allows their persistence in a variety of ecological niches and hosts [17, 20, 23, 33, 47, 93]. Infectious species and serotypes of *Vibrio* can deleteriously affect a broad range of marine taxa causing mass mortality events [18]. *Vibrio* may persist in the water column, although greater concentrations of some pathogenic species of *Vibrio* have been observed in sediment, zooplankton, mussels, and fish [19, 21, 43, 45, 55, 78]. In humans, enteric *Vibrio* infections are acquired by ingesting water or raw seafood that is contaminated with virulent or pathogenic strains of *Vibrio*. Ingesting pathogenic *Vibrio* may lead to gastroenteritis, dehydration, septicemia and in some cases death in human and experimental hosts [7, 18, 78]. Given their

mammalian physiology, marine mammals may be similarly affected following a *Vibrio* infection.

Vibrio species have been detected in marine mammals suffering from enteritis and septicemia [58, 83]. Species of *Vibrio* also were detected in harbor seals, although the virulence of these strains was unknown [32]. Measuring the abundance, distribution, and virulence potential of *Vibrio* from seals and corresponding environmental conditions may aid in identifying processes that promote pathogen proliferation and thus may impact seal health [58, 83]. The goal of this study was to investigate the dynamics and virulence potential of *Vibrio* among free-ranging and stranded Pacific harbor seals (*P. v. richardii*) off the coast of California to identify risks associated with the presence of *Vibrio*. We determined the temporal and spatial prevalence of *Vibrio* spp. in seals and examined demographic risk factors (age, sex, and body condition) and environmental conditions (precipitation, nutrients, temperature, pH, salinity, and turbidity) associated with *Vibrio* detection. *Vibrio* prevalence and species distribution in free-ranging seals were compared with those of stranded seals to better understand the role that *Vibrio* may play in the health of harbor seals. Lastly, the virulence potential of *V. parahaemolyticus* and *V. cholerae* isolates was determined by screening for virulence genes, and clinical signs associated with potentially virulent *Vibrio* were examined.

Materials and Methods

Isolation and characterization of Vibrio spp. from free-ranging and stranded seals

Free-ranging harbor seals ($n = 220$, Fig. 1) were captured on mud flats, sand spits, or rocky outcrops in San Francisco Bay (SFB), Elkhorn Slough (ES), Tomales Bay (TB) and Humboldt Bay (HB) using beach seine, drift net, or hand-held salmon nets [32]. Seals in SFB and TB were sampled during the dry (May to October) and wet (November to April) seasons from May 2010 to June 2011. Seals in ES were sampled from August to December 2010, and those in HB were sampled in June of 2011. Free-ranging seals were not sampled during the pupping season (March to April) to avoid disturbance of mother and pup pairs. Seals were weighed (± 1 kg) and restrained physically and chemically with 5mg/ml of diazepam (Hospira, Inc., Lake Forest, Illinois USA) at a dose of 0.25mg/kg. Standard length (± 1 cm) and axillary girth (± 1 cm) also were measured. Sex and age class were determined using external characteristics, mass, standard length criteria, and time of year [9, 32].




Figure 1. Sampling sites and sample sizes for free-ranging harbor seals ($n = 319$, black dots) and area between San Luis Obispo and Mendocino (black bars) where stranded animals were sampled ($n = 189$) during admission to The Marine Mammal Center in 2007 through 2011.

Harbor seals that stranded between San Luis Obispo and Mendocino were opportunistically sampled during admission for rehabilitation from May 2010 to July 2011 (Fig. 1). These individuals were primarily pups and yearlings that beached alive ($n = 53$) and appeared ill, emaciated, or were recently deceased ($n = 4$, died in transit to the rehabilitation facility, or sampled less than six hours post-mortem). Mass (± 1 kg), standard length (± 1 cm), axillary girth (± 1 cm), sex, and age class were noted as described above. Locations of stranded seals were assigned to the nearest capture location, Monterey Bay (MB, $-121.95^{\circ}\text{W} / 36.61^{\circ}\text{N}$ to $-122.11^{\circ}\text{W} / 36.95^{\circ}\text{N}$), SFB (-

122.45°W / 37.49°N to -122.80°W / 37.99°N), and TB (-122.98°W / 38.08°N to -123.09°W / 38.39°N), to compare with free-ranging seals.

Fecal samples were collected from each seal using sterile double-tip cotton swabs in Amies transport media (HealthLink, Inc. USA) inserted into the rectum of each seal. Swab samples were placed on ice, and transported to the University of California, Davis Veterinary Medical Teaching Hospital Microbiology for sample processing. One swab from each seal was placed in alkaline peptone water (Hardy Diagnostics USA) incubated at 35°C overnight in an oxygenated incubator. Plates of thiosulfate citrate bile salts agar (TCBS, Hardy Diagnostics USA), selective for *Vibrio* growth, were inoculated with the enriched swab and incubated at 35°C overnight. Individual green or yellow colonies on TCBS agar were subjected to biochemical testing to confirm genus using triple sugar iron agar slants, Christensen's urease agar slants, spot indole test, and cytochrome oxidase test. Species identification was determined using API 20E test strips (API 20E Test Kit, bioMereux, Inc., Hazelwood, MO USA). Isolates from 2010 to 2011 were cryogenically preserved (MicroBank vials, Copan Diagnostics USA) and stored at -80°C. Isolates that were not identified to the species level using biochemical testing were genetically characterized by targeting the species-specific *ToxR* gene region using the polymerase chain reaction (PCR) amplification and amplicon visualization methodology of Bauer and Rorvik (2007) [6]. If species could not be determined, or more than one species of *Vibrio* was isolated from an individual seal, the sample was categorized as *Vibrio* mixed culture for data analysis purposes. *Vibrio* prevalence and species distribution data in free-ranging seals were compared with stranded seals among locations for all years sampled.

Banked isolates from 2010 and 2011 characterized as *V. parahaemolyticus* and *V. cholerae* were screened for genes encoding virulence factors. Multiplex PCR amplification procedures previously outlined were followed with minimal modification [6, 8, 77]. Crude deoxyribonucleic acid (DNA) was extracted from each isolate using the boiling method [56], and a 1:10 dilution was used as a template for multiplex PCR amplification. Primer dilutions (Invitrogen, Inc USA), reaction buffer concentrations (dNTP Mix, GeneAmp UK; Hot Start Taq Kit, Qiagen USA), and PCR (Thermocycler, Eppendorf, USA) amplification conditions follow those of previous studies listed above. Thermostable direct hemolysin (*tdh*; 269 bp), related thermostable direct hemolysin (*trh*; 500 bp), and thermolabile hemolysin (*tl*; 450 bp) gene regions were targeted for *V. parahaemolyticus* isolates, whereas cholera toxin (*ctx*; 617 bp) and toxin co-regulated pilus (*tcp*; 385 bp) gene regions were targeted for *V. cholerae* isolates. Amplicons were separated via gel electrophoresis and visualized using 7 µl of GelStar nucleic acid stain (Gel Star, Lonza, ME USA) per 100mL of 1.5% agarose (USB Corporation, OH USA) in tri-acetate-EDTA (TAE) buffer (Bio-Rad, Inc., USA). A subset ($n = 7$) of positive samples and sample controls were sequenced (ElimBio, CA USA) to confirm amplification of target regions (Geneious 5.5, NZ). Virulence gene profiles from isolates collected from free-ranging seals were compared with stranded seals among locations. Lastly, clinical signs in seals carrying potentially virulent *Vibrio* were also documented.

Environmental data collection

Water quality and weather data collected at a resolution of 15-minute intervals were downloaded from NOAA's National Estuarine Research Reserve System (NERRS) database from SFB (SCQC1, 38.21° N / -122.03°W; SFX1, 38.22°N / -122.03°W) and ES (ELQC1 and ELXC1, 36.82°N / -121.74°W). Daily averaged salinity (ppt), temperature (°C), turbidity (NTU), pH (standard units), nutrient data (NO₃- uM, available only for ES), and daily cumulative precipitation (mm) were computed from raw, quality assured, and quality controlled NERRS data files (Matlab R2011a, Mathworks, Inc., USA) for both sampling locations. Missing or flagged data from NERRS quality control and assurance checks were excluded from analyses [94].

Data analysis

Previous *Vibrio* prevalence data collected from free-ranging seals ($n = 99$) in SFB and TB (May, June, and December of 2007, and May and June of 2008) using similar methodology were included to test the effects of environmental and demographic predictors among locations and years [32]. Chi-squared (χ^2) tests were performed to determine if the presence or absence of *Vibrio* was dependent on year, season (wet or dry), or location (SFB or TB) in free-ranging seals. A separate χ^2 test was used to compare presence or absence of *Vibrio* in free-ranging seals from ES to that of other locations sampled in 2010 to 2011. If no significant differences were detected, data were combined for subsequent analyses. Factors that were significant were either separated by factor level, or used as covariates in a logistic regression assessing risk factors. The

magnitude of effect of categorical predictor variables used in logistic regression was determined by comparing odds ratios (OR = proportion of cases/ 1- proportion of cases for reference category). Age classes were collapsed into two main categories; pups were grouped with yearlings, and subadults with adults. Residuals from the linear regression of length versus mass were used as an indicator of body condition (BCI). A forced entry, exploratory logistic regression analysis was used to determine whether age, sex, and body condition predicted the presence or absence of *Vibrio* in free-ranging harbor seals.

Stepwise lagged correlation analysis (Matlab, Student Version R2011a, Mathworks, Inc., USA) was used to determine if *Vibrio* prevalence in free-ranging seals sampled in SFB and ES was related to rainfall events for up to twenty-two days before sampling date. Environmental predictor variables measured for SFB and ES were tested for multicollinearity and normality. Daily average nutrients (ES only), temperature, salinity, pH, and turbidity values for each sample day were assigned into categories using the bin criteria determined by the mean and/or median for SFB (Appendix A) and ES (Appendix B). *Vibrio* prevalence corresponding to each category was tested for goodness of fit (Pearson's χ^2 , or Cochran's for $df = 1$). Variables with the greatest χ^2 value and correlation coefficients less than 0.70 were retained for further analysis. A backwards stepwise logistic regression was used to assess the effect of environmental predictors on the presence or absence of *Vibrio* in free-ranging seals. Odds ratios were calculated to determine the magnitude of effect.

Similarly, previous *Vibrio* prevalence data from live ($n = 102$) and recently deceased stranded seals ($n = 30$) seals in January 2007 to September 2008 were included

to test the effects of predictors on the presence or absence of *Vibrio* [32]. Chi-squared tests were performed on categorical (year, season, location, sex) predictor variables, and t-tests were performed on continuous (admission mass, BCI) predictor variables. Only significant predictors were included in a backwards stepwise logistic regression of *Vibrio* prevalence. The overall model fit was determined using the likelihood (LR) ratio test and/or the Homer and Lemeshow (HL) test statistic [68]. Statistical analyses were performed using PSAW Statistics (19.0, IMB, USA), and statistical significance was assumed for an alpha level less than 0.05 for all analyses.

Results

Vibrio prevalence in free-ranging harbor seals

From 2007 to 2011, the overall prevalence of *Vibrio* in free-ranging seals was 29% ($n = 319$). Neither year ($\chi^2_{0.05,3} = 5.512, P = 0.138$ for years sampled in SFB, and $\chi^2_{0.05,3} = 4.643, P = 0.200$ for TB) nor season (Cochran's $\chi^2_{0.05,1} = 0.276, P = 0.599$) had an effect on presence of *Vibrio* in free ranging seals, therefore, these data were pooled within locations for further analyses. Significant differences in *Vibrio* presence were detected among locations ($\chi^2_{0.05,2} = 56.237, P < 0.001$; Fig. 2). Elkhorn Slough had the greatest proportion of seals confirmed with *Vibrio* (58.7%, $n = 46$), followed by SFB (45.0%, $n = 71$), and TB (11.5%, $n = 174$). Seals sampled in ES were ten times more likely to carry *Vibrio* (OR = 10.94), whereas seals in SFB were six times more likely (OR = 6.32) than seals in TB ($P < 0.001$). The overall model was significant (LR, $P < 0.001$; HL, $P <$

1.000), and was 75% accurate in classifying presence or absence of *Vibrio* (Table 1).

Because location significantly predicted *Vibrio* in seals, data were separated by location for further analyses.




Figure 2. Proportion of free-ranging harbor seals with *Vibrio* (seals confirmed with *Vibrio*/ total seals sampled) by year and location (SFB = San Francisco Bay, TB = Tomales Bay, ES = Elkhorn Slough). Seals were sampled from ES only in 2010. Sample sizes are indicated in parentheses.

Table 1. Results from logistic regression testing location as a predictor for *Vibrio* presence and absence in seals sampled in 2007 to 2011.

Location	<i>n</i>	<i>n</i> ₁ ^a	Prevalence (%)	<i>P</i> value	Odds Ratio (OR)	CI for OR
Location 2007-2011	291	79	30.00			
Elkhorn Slough	46	27	58.69	<0.001*	10.94	5.17- 23.15
San Francisco Bay	71	32	45.07	<0.001*	6.32	3.27-3.22
Tomales Bay ^b	174	20	11.49	<0.001*	-	-
Constant				<0.001	0.17	-

^aVibrio+, ^bReference Category

Free-ranging pups and yearlings from SFB were more likely to carry *Vibrio* (52%) compared with adults and subadults (36%), although age class was not a significant predictor of *Vibrio* when tested using logistic regression ($P = 0.26$). Body condition ($P = 0.93$) and sex ($P = 0.41$) also were not significant predictors of *Vibrio* in free-ranging seals from SFB. Demographic risk factor analysis was not performed on data from seals sampled at other locations because pups and yearlings were poorly represented (ES) or presence of *Vibrio* was rare (TB).

No relationship was detected between *Vibrio* prevalence of free-ranging seals and daily average precipitation in SFB and ES, and no lag in *Vibrio* prevalence with rainfall was detected using lagged correlation analysis. The occurrence of *Vibrio* in seals was related to pH (Cochran's $\chi^2_{0.05,1} = 3.82$, $P = 0.051$ for SFB, and $\chi^2_{0.05,1} = 5.81$, $P = 0.016$ for ES), and turbidity (Cochran's $\chi^2_{0.05,1} = 8.02$, $P = 0.005$ for SFB, and $\chi^2_{0.05,1} = 7.94$, $P = 0.005$ for ES). Salinity was not significantly related, however this parameter met the criteria to be included in logistic regression (greatest chi-squared; $r < 0.70$). Temperature was significantly correlated with nutrients ($r = 0.988$, $P < 0.01$), and salinity ($r = 0.999$, $P < 0.01$), but neither temperature nor nutrients were related to *Vibrio* presence

(Appendices A & B). Salinity, pH, and turbidity ($r < 0.70$) in SFB and ES were then selected for logistic regression analysis. The most parsimonious model (LR, $P = 0.005$; HL, $P < 1.000$) for predicting *Vibrio* in free-ranging seals from SFB included only turbidity ($P = 0.008$). *Vibrio* was four and a half times more likely to occur in harbor seals sampled in SFB when the turbidity was greater than 66 NTU with a classification accuracy of 65% (Table 2). *Vibrio* in free-ranging seals from ES was best predicted by turbidity ($P = 0.003$) and salinity ($P = 0.090$) with a classification accuracy of 72% (LR, $P = 0.002$; HL, $P < 1.000$). In contrast to SFB, seals from ES were twenty-six times more likely to carry *Vibrio* when turbidity was less than 7 NTU and four and a half times more likely when salinity measured less than 33 ppt (Table 2).

Table 2. Results from multivariate backwards stepwise logistic regression testing environmental variables as predictors of the presence or absence of *Vibrio* in free-ranging seals sampled from Elkhorn Slough in 2010 and then San Francisco Bay from 2007 to 2011.

Predictors	<i>n</i>	<i>n</i> ₁ ^a	Prevalence (%)	<i>P</i> value	Odds Ratio (OR)	CI for OR
Elkhorn Slough						
Salinity (ppt)	47	29	61.70	-	-	-
<33	21	12	50.00	0.09	4.67	0.78-28.05
≥33 ^b	26	17	65.38	-	-	-
Turbidity (NTU)	47	29	61.70	-	-	-
≤ 7	17	15	88.23	0.003	26.25	3.04-226.6
> 7 ^b	30	14	46.67	-	-	-
San Francisco Bay						
Turbidity (NTU)	71	32	45.07	-	-	-
≤ 66 ^b	26	6	23.07	-	-	-
> 66	45	26	57.78	0.008	4.51	1.49-13.64

^a*Vibrio* +, ^bReference Category

Vibrio prevalence in stranded seals

The overall prevalence of *Vibrio* in stranded seals from 2007 to 2011 was 17% ($n = 189$).

Stranded adults and subadults were dropped from further analysis because of small sample size ($n = 5$). Differences in *Vibrio* prevalence were detected among years

(Cochran's $\chi^2_{0.05,1} = 10.31$, $P = 0.016$) with the greatest percentage of seals positive for

Vibrio (33 %, $n = 43$) in 2011 (Fig. 3). There was no effect of season (Cochran's $\chi^2_{0.05,1} =$

0.08, $P = 0.782$), or sex (females 20 %, $n = 49$, males 21 %, $n = 53$; Cochran's $\chi^2_{0.05,1} =$

0.002, $P = 0.966$) for the presence or absence of *Vibrio* in stranded seals.




Figure 3. Proportion of stranded pups and yearlings admitted to The Marine Mammal Center with *Vibrio*, 2007 to 2011. Sample sizes are indicated in parentheses.

Pups and yearlings with *Vibrio* had slightly greater masses at admission ($\bar{X} = 8.9$ kg, $SE = 0.49$), and a slightly greater BCI ($\bar{X} = 0.03$, $SE = 0.22$) compared with individuals without *Vibrio* (mass: $\bar{X} = 8.5$ kg, $SE = 0.30$; BCI: $\bar{X} = -0.01$, $SE = 0.11$), however, these differences were not statistically significant (mass $t(100) = -0.613$, $P =$

0.541; BCI $t(99) = -0.174$, $P = 0.862$). Since none of the demographic factors were significantly related to *Vibrio* prevalence in stranded seals, no further analyses were performed.

Prevalence of *Vibrio* in stranded seals varied by stranding county and year; however, data were too sparse to test for statistical significance. In general, San Luis Obispo (SLO), Monterey, Marin, and Mendocino counties had the greatest percentage of stranded seals positive for *Vibrio* from 2007 to 2011, although it varied among years (Fig. 1). The greatest prevalence of *Vibrio* observed per county occurred in 2011 for stranded seals sampled from SLO (67 %, $n = 3$), followed by Monterey (42 %, $n = 12$), Marin (33 %, $n = 9$), Mendocino (33 %, $n = 9$), and San Mateo (25 %, $n = 4$). Animals that stranded in Alameda and Santa Cruz counties were negative for *Vibrio* in all study years.

Comparison of Vibrio prevalence, species distribution, and virulence profiles between free-ranging and stranded seals

Because there were very few stranded adults and subadults, only data from pups and yearlings were used to compare *Vibrio* prevalence between free-ranging and stranded seals among sample locations. *Vibrio* prevalence observed in stranded and free-ranging pups and yearlings increased from 2007 to 2011 in all locations and varied among sample year (Table 3). *Vibrio* was not detected until 2011 in free-ranging pups and yearlings sampled from TB (17 %, $n = 6$). *Vibrio* prevalence was greater in free-ranging pups and yearlings compared with stranded pups and yearlings from SFB and ES. The opposite

trend was observed for TB in 2011. All free-ranging pups and yearlings sampled from ES in 2010 were positive for *Vibrio*, whereas all stranded individuals were negative.

Table 3. Prevalence of *Vibrio* in free-ranging and stranded pups and yearlings positive for *Vibrio* per location from 2007 to 2011.

Location	<i>Vibrio</i> Prevalence % (n)			
	2007	2008	2010	2011
Tomales Bay				
Free-ranging	0 (4)	0 (12)	0 (1)	17 (6)
Stranded	0 (4)	0 (6)	100 (1)	25 (4)
San Francisco				
Free-ranging	64 (11)	47 (19)	67 (6)	0 (2)
Stranded	11 (19)	17 (23)	0 (5)	60 (5)
Monterey Bay				
Free-ranging ^a	NA	NA	100 (4)	NA
Stranded	13 (8)	0 (13)	0 (3)	57 (7)


^aSampled in Elkhorn Slough

NA = did not sample

Vibrio parahaemolyticus, *V. alginolyticus*, and *V. cholerae*, were isolated from seals, although the proportions of these species were not equal in free-ranging and stranded seals among locations (Fig. 4). Isolates from free-ranging seals in ES were predominantly *V. parahaemolyticus*, whereas greater proportions of all three *Vibrio* species were observed in seals that stranded in MB. All three *Vibrio* species also were isolated from free-ranging and stranded seals from SFB, although greater proportions of *V. parahaemolyticus* and *V. alginolyticus* were isolated than *V. cholerae* in free-ranging seals. *Vibrio cholerae* was isolated primarily from stranded pups and yearlings except it was found in one free-ranging adult from ES and one free-ranging pup from SFB. The majority of isolates from free-ranging seals sampled in TB and HB were *V. alginolyticus*, with the exception of three isolates of *V. parahaemolyticus* (TB, $n = 1$; HB, $n = 2$). One

stranded seal in TB was positive for *V. parahaemolyticus*, and one for *V. cholerae*. Three isolates (SFB, $n = 2$; ES, $n = 1$) could not be identified to species level due to mixed biochemical and genotypic results.

A.



B.




Figure 4. Proportions of *Vibrio* species isolated from (A) free-ranging harbor seals sampled in Humboldt Bay (HB, $n = 13$), Tomales Bay (TB, $n = 20$), San Francisco Bay (SFB, $n = 32$) and Elkhorn Slough (ES, $n = 28$), and (B) harbor seals stranded in TB ($n = 2$), SFB ($n = 11$), and Monterey Bay (MB, $n = 6$). Mixed *Vibrio* Culture are isolates that could not be identified to species level, and samples with more than one species of *Vibrio* detected.

V. parahaemolyticus and *V. parahaemolyticus*-like species (API 20E confirmed, *ToxR*-) were detected in free-ranging and stranded seals at all locations. Ninety percent of *V. parahaemolyticus* isolates ($n = 53$) contained the *tl* gene, whereas 67% contained both *trh* and *tl* genes (Table 4). Five *V. parahaemolyticus* isolates were positive for all three hemolysin genes (*tl*, *trh*, and *tdh*), and were collected from free-ranging seals sampled in SFB ($n = 2$) and ES ($n = 3$). Overall, 77% percent of *V. parahaemolyticus* ($n = 53$) isolates contained one or more virulence genes (*trh*, *tdh*, or both), with the majority of virulent isolates from free-ranging adults and subadults in SFB and ES. Two isolates of *V. parahaemolyticus* (*ToxR*+), and three isolates biochemically similar to *V. parahaemolyticus* (*ToxR*-) were lacking all three target genes. One free-ranging seal sampled from ES was carrying *V. cholerae* (*ToxR*+) and potentially virulent *V. parahaemolyticus* (*trh*+, and *tl*+) simultaneously. All *V. cholerae* (*ToxR*+) isolates ($n = 3$) collected from seals in 2010 and 2011 demonstrated hemolytic activity on blood agar media, although they were negative for *tcp* and *ctx* virulence target genes. Target amplicon identity was confirmed by comparing sequences from a subset of *V. parahaemolyticus* and *V. cholerae* isolates with reference sequences in Genbank (Appendix C).

Table 4. Virulence gene profiles for *V. parahaemolyticus* isolates collected from free-ranging seals from Elkhorn Slough (ES), San Francisco Bay (SFB), Tomales Bay (TB), and Humboldt Bay (HB), and stranded seals (TMMC) in 2010 and 2011. Sample sizes represent the number of isolates tested.

Species	<u>Target Genes</u>			<u>Location</u>				
	<i>tdh</i>	<i>trh</i>	<i>tl</i>	ES (n=27)	SFB (n=12)	TB (n=2)	HB (n=5)	TMMC (n=7)
<i>V. parahaemolyticus</i>	+	+	+	3	2	0	0	0
	-	-	-	0	2	0	0	0
	-	+	+	20	7	1	2	1
	-	-	+	4	1	0	0	2
<i>Vibrio spp. (Vp like)</i>	-	-	-	0	0	1	1	1
	-	+	+	0	0	0	2	3

Four stranded pups sampled in 2010 and 2011 were carriers of potentially virulent *V. parahaemolyticus* (*trh*+) and *V. parahaemolyticus*-like (*trh*+) isolates, and three of those seals died in treatment. Evidence of enteritis was observed during necropsy for two of the three deceased seals carrying potentially virulent *V. parahaemolyticus*, and one carrying hemolytic *V. cholerae*.

Discussion

Pacific harbor seals off the coast of California were carriers of potentially virulent isolates of *Vibrio*. All free ranging seals appeared healthy at the time of sampling, so were asymptomatic carriers, whereas stranded seals were all underweight for their age, thus could potentially have been impacted by the *Vibrio* infections. Similar species of *Vibrio* were observed among free-ranging and stranded seals, with the exception of *V. cholerae*. *Vibrio cholerae* were primarily detected in non-weaned stranded pups and may

be of greatest concern to harbor seal health. Stranded seal pups with potentially virulent *V. parahaemolyticus* and hemolytic *V. cholerae* presented symptoms of enteritis. Future clinical studies should focus on assessing the impact of virulent *Vibrio* strains on stranded seal health.

Vibrio likely contribute to the natural microbial flora of coastal areas used by harbor seals, and population dynamics of *Vibrio* may be influenced by both oceanographic and terrestrial contributions to water quality. Changes in temperature and salinity commonly relate to the seasonal abundance and distribution of *Vibrio* in the water column [17, 19, 39, 45, 52, 55]. However, turbidity and/or salinity best predicted *Vibrio* occurrence in harbor seals, although the relationship differed with location. The differences in related environmental factors suggest that *Vibrio* exposure in seals may be related to other factors not measured in this study. Furthermore, the lack of seasonal variation for *Vibrio* prevalence in seals may indicate that seals may frequent reservoirs of *Vibrio* that are decoupled from seasonally influenced environmental factors measured here [27, 55, 86].

Differing land-use practices may influence the variation in *Vibrio* prevalence observed in seals among locations. *Vibrio* prevalence in seals from TB was the least among all the locations. Rural land surrounding TB is typically used for agriculture where the dominant enterprise is livestock farming [5]. Movement patterns and habitat use of seals from TB are unknown making it difficult to make inferences about dynamics of *Vibrio* in seals sampled at this location. However, we can conclude that it is unlikely that dairy cattle near TB are a source of *Vibrio* because TB had the greatest concentration

of dairies yet the lowest prevalence of *Vibrio* compared with ES or SFB. Furthermore, no cattle sampled near ES were confirmed with *Vibrio* during years we sampled harbor seals [65].

San Francisco Bay is the largest urbanized estuary on the eastern Pacific, and is greatly impacted by industrial and residential inputs from the Sacramento and San Joaquin rivers. Compared with ES and TB, the SFB estuary is deeper on average and comprised of a larger watershed [14]. Tidal mixing and run-off from rivers, tributaries, treatment plant effluent, industry, and groundwater contribute to changes in turbidity throughout the bay. Turbidity best predicted *Vibrio* in seals from SFB. Bacterioplankton in SFB are generally evenly distributed, and areas of greater flow near the delta facilitate conditions of optimal bacterial growth [41, 42]. No differences were detected for *Vibrio* prevalence in seals sampled from north and south SFB. Freshwater run-off from the delta also may facilitate optimal conditions for *Vibrio* growth in SFB. The turbidity maximum zone of SFB where freshwater inputs from the delta meet seawater is dominated by aggregations of particle-associated bacteria and plankton [41, 42]. *Vibrio* can associate with plankton [45, 55, 86], therefore blooms following nutrient loading from the delta may relate to the abundance of *Vibrio* in SFB. It is unknown whether *Vibrio* in SFB are free-living, or form aggregations with plankton or particulate matter in the sediment. However, if *Vibrio* aggregate near the maximum turbidity zone in SFB, seals ingesting contaminated prey or sediment in these areas may acquire greater concentrations of *Vibrio*.

Environmental dynamics relating to *Vibrio* prevalence in free-ranging seals from ES were opposite to those of SFB. It is unknown whether ES is a source for *Vibrio* in seals, yet free-ranging seals in ES had the greatest prevalence of virulent *Vibrio* overall. If ES is a source of *Vibrio* for seals, conditions that are unique to ES may relate to the observed differences. Water quality and hydrography in ES have been altered by intensive agriculture cultivation and dairy farming [14, 15]. Pesticides are continuously used for crop production and are introduced into ES from irrigation run-off, erosion, and groundwater [70, 72]. The effects of these inputs likely are magnified due to the small size and shallow bathymetry of ES compared with SFB and TB [14]. Less turbid and less saline waters were related to *Vibrio* occurrence in seals of ES. Previous studies indicated the greatest concentrations of bacteria in ES were observed near areas with the greatest freshwater inputs [72]. Tidal relaxation events coupled with continuous freshwater inputs from crop irrigation and groundwater seepage may be the driving factor promoting *Vibrio* proliferation in this location. If conditions in ES allow *Vibrio* populations to flourish, this small estuary could act as a reservoir for dense aggregations of these bacteria. Furthermore, zooplankton or fish that occur in ES may acquire *Vibrio* and become vectors when recruiting to habitat in MB [76, 86]. Data from seals in ES were limited to one sample year, therefore it is possible the greater *Vibrio* prevalence observed in these seals was an anomaly.

Alternatively, the different relationships between environmental factors and *Vibrio* prevalence in seals in ES and SFB may be explained by differences in habitat use and foraging behavior. Harbor seals that haul-out in ES generally use this habitat to rest

after foraging bouts [23, 64, 85]. Seals sampled in ES may forage in ES, although they spend the majority of their time exploiting habitat and foraging on benthic and pelagic prey (e.g. octopus, flatfish, and cusk-eel) in MB [67]. *Vibrio parahaemolyticus* was detected most often in seals from ES, and can be pathogenic in fish and humans. *Vibrio parahaemolyticus* was shown to occur less frequently than other species of *Vibrio* from water and invertebrates sampled from ES [59], and may persist in greater concentrations in sediment, offshore zooplankton, and fish [19, 45, 55, 86]. Therefore, resources and habitat in MB may be a source of greater concentrations of *Vibrio parahaemolyticus*. Depending on the source of *Vibrio* in seals from ES, environmental data from ES may not represent *Vibrio* dynamics in seals sampled here. Further research is needed to determine inter-annual variability, population dynamics, and host-vector interactions of *Vibrio* in ES and MB.

Similarly, *V. parahaemolyticus* was observed most often in seals from SFB, although seals in SFB primarily forage on benthic prey (e.g. gobies, staghorn sculpin, plainfin midshipman) within the estuary. They also forage offshore on pelagic schooling fish like northern anchovy, although these prey comprise a smaller portion of their diet [28, 62, 84]. Given *V. parahaemolyticus* can associate with sediment and zooplankton, transmission in seals from SFB may occur while foraging on benthic prey. Although total *Vibrio* prevalence was greater for seals from ES, similar proportions of *Vibrio parahaemolyticus* were observed in seals from SFB and ES. Seals from ES spend majority of their time exploiting habitat and resources in MB [23, 67, 85] therefore the

offshore life cycle of *Vibrio parahaemolyticus* may be important component for *Vibrio* dynamics in seals.

In general, *Vibrio* prevalence in free-ranging seals was greater than that in stranded seals. If *Vibrio* were acquired by ingesting contaminated prey or sediments while foraging, differences in foraging behavior between weaned and non-weaned pups may explain the observed differences in *Vibrio* prevalence. The majority of stranded seals were non-weaned pups that were abandoned or separated from their mothers that are dependent on mother's milk [32]. Stranded pups that were weaned may have foraged, although, not as successfully as healthy individuals. Weaned pups also may consume different types of prey in lesser quantities compared with older seals because of limited foraging experience and diving capacity [71]. Stranded individuals may suffer from other ailments or trauma, thereby, further hindering their abilities to successfully forage [50].

Free-ranging pups and yearlings may be at greater risk of acquiring *Vibrio* if they frequent habitat and resources in ES and SFB with greater *Vibrio* burden because they have greater home ranges than adults [50, 62, 64]. Age was a poor predictor of *Vibrio* prevalence for seals in SFB, however, the transient behavior of pups and yearlings may explain the greater *Vibrio* prevalence observed when compared with adults and subadults. If resources and space are limited near preferred habitat, juveniles may forage in alternative habitats to avoid competition with adults [25]. Transient pups and yearlings also may transmit *Vibrio* among locations if they are shedding the bacteria. Age could not be tested as a predictor of *Vibrio* in seals from ES or TB because the majority of free-ranging seals sampled in ES and TB were adults.

When comparing pups and yearlings, a greater *Vibrio* prevalence was observed in free-ranging seals than in stranded seals. All free-ranging pups and yearlings sampled were weaned, in better body condition, and likely were ingesting contaminated prey or sediment while foraging. In general, stranded weanlings and yearlings with *Vibrio* had greater body condition indices than those without. A few non-weaned and stranded pups in poor body condition were observed with *V. parahaemolyticus* and *V. cholerae*. The etiology of *Vibrio* in seals was not examined, and alternative transmission routes should be considered. Transmission also may occur via gestation, lactation, or contact with other seals shedding *Vibrio* at haul-out sites [90]. Additionally, *Vibrio* prevalence varied among locations and years for free-ranging and stranded pups and this was likely a result of poor sample sizes, or, the transient behavior of pups and yearlings. Prey abundance and distribution also may be a source of variation if pups and yearlings leave natal haul-out sites to locate prey. Regardless, it is apparent from our data that *Vibrio* prevalence in stranded pups and yearlings is increasing. This increase may be due to increased susceptibility to *Vibrio* acquisition or mother-to-pup transmission before separation. It is unlikely that the increase in prevalence reflects increased ability to culture the organisms during the study, as all isolations were performed at diagnostic laboratory with consistent practices.

Free-ranging and stranded seals not only differed by prevalence, but also the species of *Vibrio*. The proportions of *Vibrio* species differed among locations, and general trends between free-ranging and stranded seals were similar with the exception of *V. cholerae*. *Vibrio cholerae* was detected almost exclusively in non-weaned stranded

pups, and may be of greatest concern to harbor seal health. *Vibrio cholerae* are freshwater tolerant compared with other species of *Vibrio* and some strains are highly pathogenic in mammalian hosts [18, 24]. It is possible that stranded individuals may not have enough energy reserves to leave the shore and may accidentally ingest *V. cholerae* contaminated water or sediment near freshwater run-off sites. Sea otters (*Enhydra lutris*) using similar habitat consumed invertebrates near freshwater run-off sites and had similar prevalence of *V. cholerae* as stranded seals in this study [58]. Additionally, different species of *Vibrio* may associate with different prey or substrate types among regions [20, 45, 66]. This also may explain the differences in the proportions of species observed among locations for free-ranging and stranded seals.

Potentially virulent *V. parahaemolyticus* and hemolytic *V. cholerae* were observed in seals from ES and SFB. Direct mechanisms that contribute to virulence of *Vibrio* in seals have yet to be identified, although it is possible that virulence relates to conditions that are location specific [22]. Virulence expression may occur in response to environmental stressors, although few researchers have adequately tested this hypothesis [57, 79]. For example, iron is an important factor for *Vibrio* growth [80], therefore, hemolytic activity may be an adaptation selected for iron-limited oceanographic conditions. This hemolytic stress response may relate to increased virulence in the host however, *in situ* research is needed to test this hypothesis [80, 89].

The virulence gene regions targeted in this study can be up-regulated in human epidemic strains following experimental manipulation [22]. Environmental stressors introduced in *Vibrio* cell culture (e.g. pH, temperature, salinity, bicarbonate) can up-

regulate toxin production [1, 87]. Virulence expression and adhesion also increased in experimental hosts when isolates were subjected to certain growth conditions before infection [79]. Industrial contaminants and pesticides also may induce a virulence related stress response for some species of *Vibrio*, although this has not been tested *in situ* [26]. Seals suffering from contaminant burden also may become immuno-compromised, therefore, more susceptible to virulent pathogens [53, 61].

Vibrio alginolyticus was detected infrequently in free-ranging and stranded seals from all sample locations. This species is considered a pathogen of invertebrates, although it is rarely associated with disease in mammals [58]. In some cases, *V. alginolyticus* can have deleterious effects on mammalian hosts [7], although its role in the health of marine mammals is unknown. Given the low prevalence of *V. alginolyticus*, this species of *Vibrio* likely has the least impact on the health of harbor seals.

Environmental conditions resulting from climate variability may alter *Vibrio* ecology [17, 39, 49, 66, 75]. Because of this, real-time data on pathogen and host interactions relating to environmental perturbation are needed to better understand what induces virulence in the marine environment [57, 80, 88]. Furthermore, it is imperative to identify risks of potential pathogens like *Vibrio* to the health of marine mammals. In this study, we demonstrated that seals using habitat and resources near urbanized watersheds ES and SFB may have the greatest risk of acquiring potentially virulent *Vibrio*. Considering that humans share the environment and food resources with seals, potentially virulent *Vibrio* observed in seals also may be of concern to human health. Impaired watersheds like ES and SFB may be further perturbed as human populations increase

along the coast [43], and *Vibrio* may serve as a bioindicator for monitoring changes to regional ecosystem stability. It is critical to identify mechanisms of pathogen proliferation and associated risks of infection so we can forecast how aquatic pathogens may impact the health of marine mammals and the ecosystem they inhabit.

References

1. Abuaita BH, Withey JH (2009) Bicarbonate induces *Vibrio cholerae* virulence gene expression by enhancing *ToxT* activity. *Infection and Immunity* 77(9): 4111-4120.
2. Allen S (1988) Movement and activity patterns of harbor seals at Point Reyes Peninsula, California. Master's Thesis, University of California Berkeley.
3. Allen SG, Huber HR, Ribic CA et al (1989) Population dynamics of harbor seals in the Gulf of the Farallones, California. *California Fish and Game* 75(4): 224-232.
4. Azam F, Malfatti F (2007) Microbial structuring of marine ecosystems. *Nature Reviews Microbiology* 5: 782-91.
5. Barry S, King WLS (2010) Opportunities to sustain “greener” farming : comparing impacts of water quality regulations in two catchments : Lake Taupo (NZ) and Tomales Bay, California (USA). *Proceedings of the New Zealand Grassland Association* 72: 17-22.
6. Bauer A, Rørvik LM (2007) A novel multiplex PCR for the identification of *Vibrio parahaemolyticus*, *Vibrio cholerae* and *Vibrio vulnificus*. *Letters in Applied Microbiology* 45: 371-5.
7. Belkin S, Colwell R (2006) *Oceans and Health: Pathogens in the Marine Environment*. Springer Science Business Media, Inc. New York.
8. Bej AK, Patterson DP, Brasher CW et al (1999) Detection of total and hemolysin-producing *Vibrio parahaemolyticus* in shellfish using multiplex PCR amplification of *tl*, *tdh* and *trh*. *Journal of Microbiological Methods* 36: 215-225.
9. Bigg MA (1969) The harbor seal in British Columbia. *Bulletin of the Fisheries Research Board of Canada*. 172: 33.
10. Bogomolni A, Gast RJ, Ellis JC et al (2008) Victims or vectors: a survey of marine vertebrate zoonoses from coastal waters of the Northwest Atlantic. *Diseases of Aquatic Organisms* 81:13-38.
11. Bossart GD (2011) Marine mammals as sentinel species for oceans and human health. *Veterinary Pathology* 48: 676-90.
12. Brookens TJ, Harvey JT, O'Hara TM (2007) Trace element concentrations in the Pacific harbor seal (*Phoca vitulina richardii*) in central and northern California. *The Science of the Total Environment* 372: 676-92.

13. Burek KA, Gulland FMD, O'Hara TM (2008) Effects of climate change on Arctic marine mammal health. *Ecological Applications* 18: S126-34.
14. Caffrey J, Zabin C, Silberstein M et al. (2002) Introduction. In: Caffrey J, Brown M, Tyler WB et al (eds). *Changes in a California estuary; a profile of Elkhorn Slough*. Elkhorn Slough Foundation, Moss Landing, pp 1-14.
15. Caffery J, Broenkaw W (2002) Hydrography. In: Caffrey J, Brown M, Tyler WB et al (eds). *Changes in a California estuary; a profile of Elkhorn Slough*. Elkhorn Slough Foundation, Moss Landing, pp 29-42.
16. Colegrove KM, Greig DJ, Gulland FMD (2005) Causes of live strandings of northern elephant seals (*Mirounga angustirostris*) and Pacific harbor seals (*Phoca vitulina*) along the central California coast, 1992-2001. *Aquatic Mammals* 31:1-10.
17. Colwell RR (1996) Global climate and infectious disease: the cholera paradigm. *Science* 274:2025-2031.
18. Colwell RR (2006) A global and historical perspective of the genus *Vibrio*. In: Thompson FL, Austin B, Swings J (eds) *The Biology of Vibrios*, ASM Press, Washington DC, pp 3-11.
19. Deter J, Solen L, Antoine V et al (2010) Ecology of pathogenic and non-pathogenic *Vibrio parahaemolyticus* on the French Atlantic coast. Effects of temperature, salinity, turbidity and chlorophyll a. *Environmental Microbiology* 12: 929–37.
20. DePaola A, Capers GM, Alexander D (1994) Densities of *Vibrio vulnificus* in the intestines of fish from the U.S. Gulf Coast. *Applied and Environmental Microbiology* 60(3): 984–8.
21. DePaola A, Ulaszek J, Kaysner CA (2003) Molecular, serological, and virulence characteristics of *Vibrio parahaemolyticus* isolated from environmental, food and clinical sources in North America and Asia. *Applied and Environmental Microbiology* 69(7): 3999-4005.
22. DiRita VJ, Engleberg C, Heath A et al (2000) Virulence gene regulation inside and outside. *Philos Trans R Soc Lond B Bio Sci* 355: 657-665.
23. Eguchi T, Harvey JT (2005) Diving behavior of the Pacific harbor seal (*Phoca vitulina richardii*) in Monterey Bay, California. *Marine Mammal Science* 21(2): 283-295.
24. Faruque SM, Nair GB (2006) Epidemiology. In: Thompson FL, Austin B, Swings J (eds) *The biology of Vibrios*, ASM Press, Washington DC.

25. Field IC, Bradshaw CJA, Hoff J et al (2006) Age-related shifts in the diet composition of southern elephant seals expand overall foraging niche. *Marine Biology* 150: 1441-1452.
26. Ford TE (2000) Response of marine microbial communities to anthropogenic stress. *Journal of Aquatic Ecosystem Stress and Recovery* 7:75-89.
27. Fries JS, Characklis GW, Noble RT (2008) Sediment-water exchange of *Vibrio* sp. and fecal indicator bacteria: implications for persistence and transport in the Neuse River Estuary, North Carolina, USA. *Water Research* 42: 941-50.
28. Gible C (2011) Food habits of Pacific harbor seals (*Phoca vitulina richardii*) in San Francisco Bay, California. Master's Thesis, Moss Landing Marine Laboratories, San Jose State University.
29. Gilliss D, Cronquist A, Cartter M et al (2010) Morbidity and mortality weekly report. FoodNet, CDC.
http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6022a5.htm?s_cid=mm6022a5_w Accessed 30 June 2011.
30. Greig DJ (2002) Pregnancy and parturition of harbor seals in Monterey Bay, California. Master's Thesis, Moss Landing Marine Laboratories, San Jose State University.
31. Greig DJ, Gulland FMD, Rios CA, Hall AJ (2010) Hematology and serum chemistry in stranded and wild-caught harbor seals in central California: reference intervals, predictors of survival, and parameters affecting blood variables. *Journal of Wildlife Diseases* 46: 1172-84.
32. Greig DJ (2011) Health, disease, mortality and survival in wild and rehabilitated harbor seals (*Phoca vitulina*) in San Francisco Bay and along the central California coast. Dissertation, University of St. Andrews, UK <http://research-repository.st-andrews.ac.uk/handle/10023/1885>.
33. Greig DJ, Ylitalo GM, Wheeler EA et al (2011) Geography and stage of development affect persistent organic pollutants in stranded and wild-caught harbor seal pups from central California. *The Science of the Total Environment* 409: 3537-47.
34. Grimes DJ, Johnson CN, Dillon KS et al (2009) What genomic sequence information has revealed about *Vibrio* ecology in the ocean-a review. *Microbial Ecology* 58: 447-460.

35. Gulland FMD, Hall AJ (2007) Is Marine Mammal Health Deteriorating? Trends in the Global Reporting of Marine Mammal Disease. *EcoHealth* 4: 135-150.
36. Hall AJ, Frame E (2010) Evidence of domoic acid exposure in harbour seals from Scotland: a potential factor in the decline in abundance?. *Harmful Algae* 9: 489-493.
37. Hanni KD, Mazet JAK, Gulland FMD et al (2003) Clinical pathology and assessment of pathogen exposure in southern and Alaskan sea otters. *Journal of Wildlife Diseases* 39: 837-50.
38. Härkönen T, Dietz R, Reijnders P et al (2006) The 1988 and 2002 phocine distemper virus epidemics in European harbour seals. *Diseases of Aquatic Organisms* 68: 115-130.
39. Harvell CD, Mitchell CE, Wad JR et al (2002) Climate warming and disease risks for terrestrial and marine biota. *Science* 296: 2158-62.
40. Hashizume M, Faruque ASG, Wagatsuma Y et al (2010) Cholera in Bangladesh: climatic components of seasonal variation. *Epidemiology* 21(5):706-710.
41. Hollibaugh JT, Wong PS (1996) Distribution and activity of bacterioplankton in San Francisco Bay. In: Hollibaugh JT (ed) *San Francisco Bay the ecosystem; further investigations into the natural history of San Francisco Bay and delta with reference to the influence of man*. Pacific Division of the American Association for the Advancement of Science, San Francisco, pp 263-288.
42. Hollibaugh JT, Wong PS (1999) Microbial processes in the San Francisco Bay estuarine turbidity maximum. *Estuaries* 22(4): 848-862.
43. Hsieh JL, Fries JS, Noble RT (2008) Dynamics and predictive modelling of *Vibrio* spp. in the Neuse River Estuary, North Carolina, USA. *Environmental Microbiology* 10: 57-64.
44. Hove-Musekwa SD, Nyabadza F, Chiyaka C et al (2011) Modelling and analysis of the effects of malnutrition in the spread of cholera. *Mathematical and Computer Modelling* 53: 1583–1595.
45. Kaneko T, Colwell RR (1973) Ecology of *Vibrio parahaemolyticus* in Chesapeake Bay. *Journal of Bacteriology* 113: 24–32.
46. Keasler SP, Hall RH (1993) Detecting and biotyping *Vibrio cholerae* O1 with multiplex polymerase chain reaction. *Lancet* 341:1661.

47. Kirkup BC, Chang L, Chang S et al (2010) *Vibrio* chromosomes share common history. BMC Microbiology 10:137.
48. Knap A, Dewailly É, Furgal C, Galvin J, et al (2002) Indicators of ocean health and human health: developing a research and monitoring framework. Environmental Health Perspectives 110:839–845.
49. Lafferty KD (2009) The ecology of climate change and infectious diseases. Ecology 90: 888-900.
50. Lander ME (1998) Success of free-ranging and rehabilitated harbor seal (*Phoca vitulina richardii*) pups in the wild. Master's Thesis. Moss Landing Marine Laboratories, San Francisco State University.
51. Laws EA, Fleming LE, Stegeman JJ (2008) Centers for Oceans and Human Health: contributions to an emerging discipline. Environmental Health 7 Supplement.
52. Lipp EK, Huq A, Colwell RR (2002) Effects of global climate on infectious disease : the cholera model. Clinical Microbiology Reviews 15: 757-770.
53. Loveren VH, Ross PS, Osterhaus AD, Vos JG (2000) Contaminant-induced immunosuppression and mass mortalities among harbor seals. Toxicology Letters 112-113: 319-324.
54. Lowry LF, Frost KJ, Ver Hoef JM et al (2001) Movements of satellite-tagged subadult and adult harbor seals in Prince William Sound, Alaska. Marine Mammal Science 17: 835-861.
55. Martinez-Urtaza J, Blanco-Abad V, Rodriguez-Castro A, et al (2012) Ecological determinants of the occurrence and dynamics of *Vibrio parahaemolyticus* in offshore areas. ISME J 6(5): 994-1006.
56. Medici DD, Croci L, Delibato E et al (2003) Evaluation of DNA extraction methods for use in combination with SYBR green I real-time PCR to detect *Salmonella enterica* serotype enteritidis in poultry. Applied and Environmental Microbiology 69: 3456-346.
57. Mekalanos JJ (1992) Environmental signals controlling expression of virulence determinants in bacteria. Journal of Bacteriology 174: 1-7.
58. Miller MA, Byrne BA, Jang SS et al (2010) Enteric bacterial pathogen detection in southern sea otters (*Enhydra lutri nereis*) is associated with coastal urbanization and freshwater run-off. Veterinary Research 41: 1-13.

59. Miller WA, Miller MA, Gardner IA et al (2006) *Salmonella* spp., *Vibrio* spp. *Clostridium perfringens*, and *Plesiomonas shigelloides* in marine and freshwater invertebrates from coastal California ecosystems. *Microbial Ecology* 52(2):198-206.
60. Moore S (2008) Marine mammals as ecosystem sentinels. *Journal of Mammalogy* 89: 534-540.
61. Neale JCC, Van de Water JA, Harvey JT, Tjeerdema RS, Gershwin ME (2002) Proliferative responses of harbor seal (*Phoca vitulina*) T lymphocytes to model marine pollutants. *Developmental Immunology* 9: 215-221.
62. Nickel B (2003) Movement and habitat use patterns of harbor seals in the San Francisco estuary, California. Master's Thesis, Moss Landing Marine Laboratories, San Francisco State University.
63. Nigro OD, Hou A, Vithanage G, Fujioka RS, Steward GF (2011) Temporal and spatial variability in culturable pathogenic *Vibrio* spp. in Lake Pontchartrain, Louisiana, following hurricanes Katrina and Rita. *Applied and Environmental Microbiology* 77: 5384-5393.
64. Oates S (2005) Survival, movements, and diet of juvenile harbor seals along central California. Master's Thesis, Moss Landing Marine Laboratories, San Jose State University.
65. Oates SC, Miller MA, Byrne BA et al. (2012) Epidemiology and potential land-sea transfer of enteric bacteria from terrestrial to marine species in the Monterey Bay region of California. *Journal of Wildlife Diseases* 48(3): 654-668.
66. Ottaviani D, Leoni F, Rocchegiani et al (2012) An extensive investigation into the prevalence and the genetic and serological diversity of toxigenic *Vibrio parahaemolyticus* in Italian marine coastal waters. *Environmental Microbiology* doi: 10.1111/j.1462-2920.2012.02839.x
67. Oxman D (1995) Seasonal abundance, movements, and food habits of harbor seals (*Phoca vitulina richardii*) in Elkhorn Slough, California. Master's Thesis, Moss Landing Marine Laboratories, California State University, Fresno
68. Peng CYJ, Lee KL, Ingersoll GM (2002) An introduction to logistic regression analysis and reporting. *The Journal of Educational Research* 96: 3-14.
69. Perrin WF, Wursig B, Thewissen JGM (2002) *Encyclopedia of marine mammals*. Elsevier, Burlington.

70. Phillips B, Stephenson M, Jacobi M, et al. (2002) Land use and contaminants. In: Caffrey J, Brown M, Tyler WB et al (eds). Changes in a California estuary; a profile of Elkhorn Slough. Elkhorn Slough Foundation, Moss Landing, pp 237-256.
71. Prewitt JS, Freistroffer DV, Schreer JF, Hammill MO, Burns JM (2010) Postnatal development of muscle biochemistry in nursing harbor seal (*Phoca vitulina*) pups: limitations to diving behavior? *Journal of Comparative Physiology B* 180: 757-766.
72. Rice DW, Seltenrich CP, Spies RB, Keller ML (1993) Seasonal and annual distribution of organic contaminants in marine sediments from Elkhorn slough, moss landing harbor and nearshore Monterey Bay, California. *Environmental Pollution* 82: 79-91.
73. Rose JM, Gast RJ, Bogolmoni A et al (2009) Occurrence and patterns of antibiotic resistance in vertebrates off the Northeastern United States coast. *FEMS Microbiology Ecology* 67: 421-431.
74. Scott TM, Rose JB, Jenkins TM, Farrah SR (2002) Microbial source tracking : current methodology and future directions. *Applied and Environmental Microbiology* 68: 5796-5803.
75. Sedas VTP (2007) Influence of environmental factors on the presence of *Vibrio cholerae* in the marine environment: a climate link. *J Infect Developing Countries* 1: 224-241.
76. Senderovich Y, Izhaki I, Halpern M (2010) Fish as reservoirs and vectors of *Vibrio cholerae*. *PloS One* 5(1): e8607.
77. Shangkuan YH, Show YS, Wang TM (1995) Multiplex polymerase chain reaction to detect toxigenic *Vibrio cholerae* and to biotype *Vibrio cholerae* O1. *The Journal of Applied Bacteriology* 79: 264-273.
78. Stewart JR, Gast RJ, Fujioka RS, Solo-Gabriele HS et al (2008) The coastal environment and human health: microbial indicators, pathogens, sentinels and reservoirs. *Environmental Health* 7(S2):S3.
79. Sung H, Chang C, Lan S (2004) Effects of salinity and pH on the adherence and virulence of *Vibrio cholerae* O139. *Journal of Food and Drug Analysis* 12(1): 68-73.

80. Sun-Hee A, Han JH, Lee JH et al (2005) Identification of an iron-regulated hemin-binding outer membrane protein, HupO, in *V. fluvialis*: effects on hemolytic activity and the oxidative stress response. *Infection and Immunity* 73(2): 722-729.
81. Tao Z, Bullard S, Arias C (2011) High numbers of *Vibrio vulnificus* in tar balls collected from oiled areas of the north-central Gulf of Mexico following the 2010 BP Deepwater Horizon Oil Spill. *EcoHealth*. doi:10.1007/s10393-011-0720-z
82. Thompson B, Adelsbach T, Brown C et al (2007) Biological effects of anthropogenic contaminants in the San Francisco Estuary. *Environmental Research* 105: 156-174.
83. Thornton SM, Nolan S, Gulland FMD (1998) Bacterial Isolates from California sea lions (*Zalophus californianus*), Pacific harbor seals (*Phoca vitulina*) and elephant seals admitted to a rehabilitation center along the central California coast. *Journal of Zoo and Wildlife Medicine* 29(2): 171-178.
84. Torok M (1994) Movements, daily activity patterns, dive behavior, and food habits of harbor seals (*Phoca vitulina richardii*) in San Francisco Bay, California. Master's Thesis, Moss Landing Marine Laboratories, San Jose State University.
85. Trumble S (1995) Abundance, movements, dive behavior, food habits, and mother-pup interactions of harbor seals near Monterey Bay, California. Master's Thesis, Moss Landing Marine Laboratories, California State University, Fresno.
86. Turner JW, Good B, Cole D, Lipp EK (2009) Plankton composition and environmental factors contribute to *Vibrio* seasonality. *ISME J* 3:1082-1092
87. Whitaker WB, Parent MA, Naughton LM et al (2010) Modulation of responses of *Vibrio parahaemolyticus* O3:K6 to pH and temperature stresses by growth at different salt concentrations. *Applied and Environmental Microbiology* 76: 4720-4729.
88. Woolhouse, M. (2011) How to make predictions about future infectious disease risks. *Philosophical Transactions of the Royal Society of London B* 366: 2045-54.
89. Wright AC, Simpson LM, Oliver JD (1981) Role of iron in the pathogenesis of *Vibrio vulnificus* infections. *Infection and Immunity* 34 (2): 503.
90. Xavier MN, Paixão TA, Hartigh ABD, Tsohis RM, Santos RL (2010) Pathogenesis of *Brucella* spp . *The Open Veterinary Science Journal* 4: 109-118.
91. Yochem PK, Stewart BS, DeLong RL, DeMaster DP (1987) Diel haul-out patterns and site fidelity of harbor seals (*Phoca vitulina richardsi*) on San Miguel Island, California, in Autumn. *Marine Mammal Science* 3: 323-332.

92. Zarnke RL, Saliki JT, Macmillan AP et al (2006) Serologic survey for *Brucella* spp., phocid herpesvirus-1, phocid herpesvirus-2, and phocine distemper virus in harbor seals from Alaska, 1976-1999. *Journal of Wildlife Diseases* 42: 290-300.
93. Zo YG, Chokesajjawatee N, Arawaka E et al (2008) Covariability of *Vibrio cholerae* microdiversity and environmental parameters. *Applied and Environmental Microbiology* 74: 2915-20.
94. _____ (2012) NOAA's National Estuarine Research Reserve System National Monitoring Program. Centralized Data Management Office, <http://cdmo.baruch.sc.edu/> Accessed February 2012.

Appendix A. Descriptive statistics, and bin criteria used for categorizing environmental predictor variables to be used in multivariate logistic regression analysis for SFB, from 2007 to 2011. Presence and absence of *Vibrio* were tested between categories for each predictor using a Pearson's chi-squared test, and associated p-values are reported.

Predictor Variable	<i>n</i> ^a	Min	Max	Mean	Median	S.E.	S.D.	Bin Criteria	Cochran's X ² , P
Temperature (*C)	873	7.19	23.88	16.51	17.22	0.13	3.92	<16, ≥16	0.640, P = 0.424
Salinity (ppt)	874	7.97	28.22	23.23	24.78	0.15	4.32	<23, ≥23	2.22, P = 0.136
pH (standard units)	837	7.65	9.08	8.03	7.98	0.01	0.20	≤8, >8	3.82, P = 0.051
Turbidity (NTU)	874	7.51	603.94	66.37	36.59	2.61	77.18	≤66, >66	8.02, P = 0.005

^aDaily values collected for sampling duration

Appendix B. Descriptive statistics, and bin criteria used for categorizing environmental predictor variables to be used in multivariate logistic regression analysis for ES in 2010. Presence and absence of *Vibrio* were tested between categories for each predictor using a Pearson's chi-squared test, and associated p-values are reported.

Predictor Variable	<i>n</i> ^a	Min	Max	Mean	Median	S.E.	S.D.	Bin Criteria	Cochran's X ² , P
Nutrients (NO ₃ - uM)	47	16.17	108.11	59.64	24.92	6.43	44.12	≤24.92, >24.92	0.01, P=0.901
Temperature (*C)	152	9.37	20.29	15.91	16.53	0.25	3.07	<16, ≥16	0.01, P=0.908
Salinity (ppt)	152	28.45	34.81	33.35	33.48	0.09	1.15	<33, ≥33	0.33, P=0.563
pH (standard units)	101	7.79	8.26	7.98	7.97	0.01	0.1	≤7.9, >7.9	5.81, P=0.016
Turbidity (NTU)	140	3.43	19.39	7.37	6.80	0.21	2.54	≤7, >7	7.94, P=0.005

^aDaily values collected for sampling duration

Appendix C. Sequence confirmation for target genes for a subset of isolates collected from harbor seals.

Harbor Seal Isolate	Target Region	Observed Length	Expected	E-Value	% Pairwise Identity	Strain & Reference Accession #
<i>V.parahaemolyticus</i>						
<i>tdh, trh, tl</i> ^a						
Adult Female C	<i>tdh</i>	313	270	5.13E-130	98.2	<i>V.parahaemolyticus</i> 03:K6 (Bangladesh), AY044114
	<i>trh</i>	482	500	0	97.6	<i>V.parahaemolyticus</i> TH3996, AB455531
	<i>tl</i>	447	450	0	98.7	<i>V.parahaemolyticus</i> , AY289609
1972-1973	<i>tdh</i>	315	270	8.94E-128	97.9	<i>V.parahaemolyticus</i> 03:K6 (Bangladesh), AY044114
	<i>trh</i>	481	500	0	98.5	<i>V.parahaemolyticus</i> , AY742213
	<i>tl</i>	434	450	0	99.1	<i>V.parahaemolyticus</i> ATCC 33846, GU971655
1935-1936	<i>tdh</i>	309	270	4.04E-131	98.5	<i>V.parahaemolyticus</i> 03:K6 (Bangladesh), AY044114
	<i>trh</i>	481	500	0	97.9	<i>V.parahaemolyticus</i> , AY742213
	<i>tl</i>	434	450	0	98.8	<i>V.parahaemolyticus</i> , AY289609
1892-1893A	<i>tdh</i>	306	270	6.70E-129	97.8	<i>V.parahaemolyticus</i> 03:K6 (Bangladesh), AY044114
	<i>trh</i>	445	500	0	98.2	<i>V.parahaemolyticus</i> , AY742213
	<i>tl</i>	436	450	0	98.7	<i>V.parahaemolyticus</i> , AY289609
<i>V.parahaemolyticus</i>						
<i>ToxR</i> ^b						
2014-2015	<i>ToxR</i>	274	297	3.59E-133	98.5	<i>V.parahaemolyticus</i> RIMD 2210086, AY527397
1822-1823C	<i>ToxR</i>	246	297	2.56E-109	96.3	<i>V.parahaemolyticus</i> RIMD 2210086, AY527397
<i>V.cholerae</i>						
<i>ToxR</i> ^b						
1824-0491	<i>ToxR</i>	573	640	0	98.6	<i>V.cholerae</i> 01 El Tor N16961, AE003852

^aBej et al. 1999, ^bBauer & Rorvik 2007